

Scotland's Rural College

Paternal Origins and Migratory Episodes of Domestic Sheep

Deng, Juan; Xie, Xing Long; Wang, Dong Feng; Zhao, Chao; Lv, Feng Hua; Li, Xin; Yang, Ji; Yu, Jia Lin; Shen, Min; Gao, Lei; Yang, Jing Quan; Liu, Ming Jun; Li, Wen Rong; Wang, Yu Tao; Wang, Feng; Li, Jin Quan; Hehua, EE E.; Liu, Yong Gang; Shen, Zhi Qiang; Ren, Yan Ling; Liu, Guang Jian; Chen, Ze Hui; Gorkhali, Neena A.; Rushdi, Hossam E.; Salehian-Dehkordi, Hosein; Esmailizadeh, Ali; Nosrati, Maryam; Paiva, Samuel R.; Caetano, Alexandre R.; Štěpánek, Ondřej; Olsaker, Ingrid; Weimann, Christina; Erhardt, Georg; Curik, Ino; Kantanen, Juha; Mwacharo, Joram M.; Hanotte, Olivier; Bruford, Michael W.; Ciani, Elena; Periasamy, Kathiravan; Amills, Marcel; Lenstra, Johannes A.; Han, Jian Lin; Zhang, Hong Ping; Li, Li; Li, Meng Hua

Published in:
Current Biology

DOI:
[10.1016/j.cub.2020.07.077](https://doi.org/10.1016/j.cub.2020.07.077)

Print publication: 19/10/2020

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for pulished version (APA):

Deng, J., Xie, X. L., Wang, D. F., Zhao, C., Lv, F. H., Li, X., Yang, J., Yu, J. L., Shen, M., Gao, L., Yang, J. Q., Liu, M. J., Li, W. R., Wang, Y. T., Wang, F., Li, J. Q., Hehua, EE. E., Liu, Y. G., Shen, Z. Q., ... Li, M. H. (2020). Paternal Origins and Migratory Episodes of Domestic Sheep. *Current Biology*, 30(20), 4085-4095.e6. <https://doi.org/10.1016/j.cub.2020.07.077>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Current Biology

Paternal Origins and Migratory Episodes of Domestic Sheep

Highlights

- Novel ovine SNPs of the male-specific region of Y chromosome were developed
- Y chromosome of domestic sheep contains four different paternal lineages
- Lineages C and B predominate in breeds of primitive traits and fat tail, respectively
- Expansions of sheep correlate with various phenotypic traits and breeding goals

Authors

Juan Deng, Xing-Long Xie,
Dong-Feng Wang, ...,
Hong-Ping Zhang, Li Li, Meng-Hua Li

Correspondence

menghua.li@cau.edu.cn

In Brief

Deng et al. show that domestic sheep harbor four Y chromosome lineages and early expansions of sheep were associated with the segregation of primitive and fat-tailed phenotypes as well as traits selected for different purposes.

Report

Paternal Origins and Migratory Episodes of Domestic Sheep

Juan Deng,^{1,2,39} Xing-Long Xie,^{1,3,39} Dong-Feng Wang,^{1,3} Chao Zhao,^{1,4} Feng-Hua Lv,^{1,5} Xin Li,^{1,3} Ji Yang,^{1,5} Jia-Lin Yu,⁶ Min Shen,^{7,8} Lei Gao,^{7,8} Jing-Quan Yang,^{7,8} Ming-Jun Liu,⁹ Wen-Rong Li,⁹ Yu-Tao Wang,¹⁰ Feng Wang,¹¹ Jin-Quan Li,¹² EEer Hehua,¹³ Yong-Gang Liu,¹⁴ Zhi-Qiang Shen,¹⁵ Yan-Ling Ren,¹⁵ Guang-Jian Liu,¹⁶ Ze-Hui Chen,^{1,3} Neena A. Gorkhali,¹⁷ Hossam E. Rushdi,¹⁸ Hosein Salehian-Dehkordi,^{1,3} Ali Esmailzadeh,¹⁹ Maryam Nosrati,²⁰

(Author list continued on next page)

¹CAS Key Laboratory of Animal Ecology and Conservation Biology, Institute of Zoology, Chinese Academy of Sciences (CAS), Beijing 100101, China

²Farm Animal Genetic Resources Exploration and Innovation Key Laboratory of Sichuan Province, Sichuan Agricultural University, Chengdu 611130, China

³University of Chinese Academy of Sciences, Beijing 100049, China

⁴College of Life Science, Hebei University, Baoding 071002, China

⁵College of Animal Science and Technology, China Agricultural University, Beijing 100193, China

⁶Station for Breeding and Improvement of Animal and Poultry of Changshou District, Chongqing 401220, China

⁷Institute of Animal Husbandry and Veterinary Medicine, Xinjiang Academy of Agricultural and Reclamation Sciences, Shihezi 832000, China

⁸State Key Laboratory of Sheep Genetic Improvement and Healthy Breeding, Xinjiang Academy of Agricultural and Reclamation Sciences, Shihezi 832000, China

⁹Animal Biotechnological Research Center, Xinjiang Academy of Animal Science, Urumqi 830001, China

¹⁰College of Life and Geographic Sciences, Kashi University, Kashi 844000, China

¹¹Institute of Sheep and Goat Science, Nanjing Agricultural University, Nanjing 210095, China

¹²College of Animal Science, Inner Mongolia Agricultural University, Hohhot 010000, China

¹³Grass-Feeding Livestock Engineering Technology Research Center, Ningxia Academy of Agriculture and Forestry Sciences, Yinchuan 750000, China

¹⁴College of Animal Science and Technology, Yunnan Agricultural University, Kunming 650000, China

¹⁵Shandong Binzhou Academy of Animal Science and Veterinary Medicine, Binzhou 256600, China

¹⁶Novogene Bioinformatics Institute, Beijing 100083, China

(Affiliations continued on next page)

SUMMARY

The domestication and subsequent global dispersal of livestock are crucial events in human history, but the migratory episodes during the history of livestock remain poorly documented [1–3]. Here, we first developed a set of 493 novel ovine SNPs of the male-specific region of Y chromosome (MSY) by genome mapping. We then conducted a comprehensive genomic analysis of Y chromosome, mitochondrial DNA, and whole-genome sequence variations in a large number of 595 rams representing 118 domestic populations across the world. We detected four different paternal lineages of domestic sheep and resolved, at the global level, their paternal origins and differentiation. In Northern European breeds, several of which have retained primitive traits (e.g., a small body size and short or thin tails), and fat-tailed sheep, we found an overrepresentation of MSY lineages y-HC and y-HB, respectively. Using an approximate Bayesian computation approach, we reconstruct the demographic expansions associated with the segregation of primitive and fat-tailed phenotypes. These results together with archaeological evidence and historical data suggested the first expansion of early domestic hair sheep and the later expansion of fat-tailed sheep occurred ~11,800–9,000 years BP and ~5,300–1,700 years BP, respectively. These findings provide important insights into the history of migration and pastoralism of sheep across the Old World, which was associated with different breeding goals during the Neolithic agricultural revolution.

RESULTS AND DISCUSSION

The Neolithic agricultural revolution triggered the transition from a hunter-gatherer nomad lifestyle to a sedentary farming society,

and a subsequent spreading of agriculture and pastoralism across Asia, Europe, and Africa during ~10,000–5,000 years before present (BP) [4, 5]. Previous molecular studies have contributed to the reconstruction of domestication and

Samuel R. Paiva,²¹ Alexandre R. Caetano,²¹ Ondřej Štěpánek,²² Ingrid Olsaker,²³ Christina Weimann,²⁴ Georg Erhardt,²⁴ Ino Curik,²⁵ Juha Kantanen,²⁶ Joram M. Mwacharo,^{27,28} Olivier Hanotte,^{29,30} Michael W. Bruford,^{31,32} Elena Ciani,³³ Kathiravan Periasamy,³⁴ Marcel Amills,³⁵ Johannes A. Lenstra,³⁶ Jian-Lin Han,^{37,38} Hong-Ping Zhang,² Li Li,² and Meng-Hua Li^{1,5,40,*}

¹⁷Animal Breeding Division, National Animal Science Institute, Nepal Agriculture Research Council (NARC), Kathmandu, Nepal

¹⁸Department of Animal Production, Faculty of Agriculture, Cairo University, 12613 Giza, Egypt

¹⁹Department of Animal Science, Faculty of Agriculture, Shahid Bahonar University of Kerman, Kerman, Iran

²⁰Department of Agriculture, Payame Noor University, Tehran, Iran

²¹Embrapa Recursos Genéticos e Biotecnologia, Parque Estação Biológica, PqEB, Avenida W5 Norte (Final), Caixa Postal 02372, CEP 70770-917 Brasília, DF, Brazil

²²Department of Virology, State Veterinary Institute Jihlava, Rantirovská 93, 58601, Jihlava, Czech Republic

²³Department of Preclinical Sciences and Pathology, Faculty of Veterinary Medicine, Norwegian University of Life Sciences, Oslo, Norway

²⁴Department of Animal Breeding and Genetics, Justus-Liebig-University Giessen, Giessen, Germany

²⁵Department of Animal Science, Faculty of Agriculture, University of Zagreb, Zagreb, Croatia

²⁶Production Systems, Natural Resources Institute Finland (Luke), FI-31600 Jokioinen, Finland

²⁷International Center for Agricultural Research in the Dry Areas (ICARDA), P.O. Box 5689, Addis Ababa, Ethiopia

²⁸CTLGH and SRUC, the Roslin Institute Building, Easter Bush Campus, Edinburgh EH25 9RG, UK

²⁹LiveGene, International Livestock Research Institute (ILRI), P.O. Box 5689, Addis Ababa, Ethiopia

³⁰School of Life Sciences, University of Nottingham, University Park, Nottingham, NG72RD, UK

³¹School of Biosciences, Cardiff University, Cathays Park, Cardiff CF10 3AX, Wales, United Kingdom

³²Sustainable Places Research Institute, Cardiff University CF10 3BA, Wales, United Kingdom

³³Dipartimento di Bioscienze, Biotecnologie e Biofarmaceutica, Università degli Studi di Bari Aldo 24 Moro, Bari, Italy

³⁴Animal Production and Health Laboratory, Joint FAO/IAEA Division, International Atomic Energy Agency (IAEA), Vienna, Austria

³⁵Department of Animal Genetics, Center for Research in Agricultural Genomics (CRAG), CSIC-IRTA-UAB-UB, Campus de la Universitat Autònoma de Barcelona, Bellaterra 08193, Spain

³⁶Faculty of Veterinary Medicine, Utrecht University, Utrecht, the Netherlands

³⁷CAAS-ILRI Joint Laboratory on Livestock and Forage Genetic Resources, Institute of Animal Science, Chinese Academy of Agricultural Sciences (CAAS), Beijing 100193, China

³⁸Livestock Genetics Program, International Livestock Research Institute (ILRI), Nairobi 00100, Kenya

³⁹These authors contributed equally

⁴⁰Lead Contact

*Correspondence: menghua.li@cau.edu.cn

<https://doi.org/10.1016/j.cub.2020.07.077>

migrations of farm animals in terms of time, place, and driving forces [1, 5–8]. Sheep was one of the first species domesticated in the Fertile Crescent ~12,000–10,000 years BP [9, 10] and initially was bred as a source of meat [11]. Domesticated sheep accompanied human migrations worldwide (Figure 1A; Data S1) and acted as a key factor in the social transformation between sedentary agrarian, seminomadic, and nomadic pastoralist societies [12–15]. In addition to the initial dispersal, archaeological records show at least other population expansions, when sheep were bred for wool and for fat tails as adaptation to a dry climate during the late Holocene (6,000–3,000 years BP) (Figure 1B; Data S1).

Autosomal and mitochondrial DNA (mtDNA) variations as well as whole-genome sequences have been widely used in revealing the domestication, migration, and genetic diversity of sheep [12, 16–18] and other farm animals [7, 8, 19, 20]. Variation of the male-specific region of the Y chromosome (MSY) can be particularly powerful because it represents the paternal lineage and does not recombine. However, it is hindered by the abundance of ampliconic regions with multicopy genes and by the heterochromatic repeat units [21, 22]. As a consequence, only few reports have studied variation in the MSY of sheep on the basis of three markers [23–26]. In this study, we used whole-genome sequences to generate a unique large set of novel MSY SNPs in ovine species. To elucidate the history of sheep domestication and their early dispersions, MSY SNP genotypes were analyzed in combination with mtDNA variations and whole-genome

sequences in a large number of rams across the world (Data S2). These include populations from Northern Europe, Eastern Europe, UK, Southern Asia, Central Asia, Mongolia, Russia, Africa, and Southern America, which were underrepresented in earlier work but essential to assess ovine population expansions. More specifically, we aim to answer several key questions: (1) how many paternal lineages exist in domestic sheep, (2) are sheep with distinct phenotypic traits and their migration episodes associated with the segregation of different lineages, and (3) when did the expansion event(s) occur?

From the high-depth whole-genome sequences of 31 rams and 5 ewes (NCBI: PRJNA624020) [27] with a wide geographic origin ranging from China and Afghanistan to the Netherlands, Sweden, UK, and France (average coverage = 24.7 × ; Data S2), we extracted a reference panel of 467 reliable single-copy MSY (scpMSY) contigs (501 kb) covering 10 Mb assembly by using a modified method in Wallner et al. [28] (Figure 2). To detect variants, we mapped whole-genome sequences of 161 rams and 32 ewes from 88 domestic populations, and 18 rams from three wild species (nine Asiatic mouflons [*O. orientalis*], eight argali [*O. ammon*], and one urial [*O. vignei*]; average coverage = 20.3 × ; NCBI: PRJNA624020 and PRJNA645671; Data S2) to the scpMSY reference. In total, we called 85 novel MSY SNPs in domestic sheep and 425 in the wild species (Data S3), 15 of which were shared by wild and domestic sheep. In the 179 domestic and wild rams, a total of 63 haplotypes were defined based on genotypes of the 495 SNPs including




Figure 1. Archaeological and Historical Records about Domestic Sheep and, in Particular, Wool and Fat-Tailed Sheep in the World

(A) Archaeological evidence for domestic sheep across the world. The dot labeled “1” means the approximate location in Central Balkans.

(B) Archaeological evidence and historical records for the wool and fat-tailed sheep. Labels 1 and 2 indicate the approximate locations of the sites (about 6,000 BC and 4,000–3,000 BC) in Iran and Central Balkans (3,300 BC), respectively. Label 3 indicates the approximate location of the wool record in Egypt. Labels 4, 5, and 6 indicate the approximate locations of the fat-tailed records in Kangju (now in Southern Kazakhstan), Hami (now in Xinjiang, China), and Mongolia, respectively.

See also [Data S1](#)

the two previously reported SNPs $\alpha Y1$ and $\alpha Y2$ [26] ([Data S4](#)). Different phylogenetic reconstruction methods yielded similar phylogenetic trees, in which two haplotypes of Asiatic mouflons are closely related to those of domestic sheep ([Figure S1A](#)). Diagnostic SNPs evidenced the segregation of four distinct haplogroups (y-HA, y-HB, y-HC, and y-HD) in domestic rams ([Figure S1A](#); [Data S4](#)), all of which were supported by high bootstrap values.

For the 179 rams, we constructed Bayesian trees based on the MSY SNPs, mitogenomes, and a neighbor-joining (NJ) tree based on whole-genome sequences ([Figure 3](#)). Two MSY haplotypes from four Asiatic mouflons (TH.2, SH.7, 266, and 267) are similar to the y-HB haplotypes from domestic sheep ([Figure S1B](#)). However, the haplotypes of five other mouflons cluster with an urial haplotype, suggesting gene flow from urial into the Asiatic mouflon population in Iran [17]. Previous field and genetic evidence have shown the presence of hybrids between urial and Asiatic mouflon in this country [29, 30]. Inter-species introgressions have also been detected in several genera [31], including

the wild and domestic sheep species [13, 32]. We observed different patterns of MSY, mtDNA, and whole-genome phylogenies among the wild and domestic sheep. Only mt-HA and mt-HB were detected in the individuals of MSY lineage y-HC; conversely, the sheep of mt-HC showed paternal lineages of y-HA and y-HB ([Figures S1B, S1C, S1E, and S1F](#)). A phylogenetic tree of the whole-genome sequences shows a significant phylogeographic pattern by clustering sheep according to their continents with the exception of six Asian fine-wool (Merino-like) sheep that are attached to the European cluster ([Figures 3C and S1D](#)).

The three major MSY haplogroups differ in Watterson θ (θ_W) values: $\theta_W = 2.47 \times 10^{-5}$ for y-HB, 2.32×10^{-5} for y-HA, and 1.04×10^{-5} for y-HC, about 200 times lower than those found for the three major

mtDNA lineages of the same samples (mt-HA, $\theta_W = 4.39 \times 10^{-3}$; mt-HB, 7.17×10^{-3} ; mt-HC, 4.19×10^{-3}). Our estimates of θ_W for MSY haplogroups are similar to those reported for domestic Bactrian camels ($\theta_W = 1.17 \times 10^{-5}$) [33] and domestic stallions ($\theta_W = 0.79 \times 10^{-5}$) [28]. Using an empirical Bayesian approach [34], we detected more recent estimates of divergence times among MSY haplogroups than for mtDNA lineages from the same individuals ([Figures 3A and 3B](#)).

Using the T_{MRCA} (time to the most recent common ancestor) of argali and urial with domestic sheep, we estimate that the overall mutation rate is 0.93×10^{-10} mutations per generation per site [95% highest posterior density interval (95% HPDI): 0.83×10^{-10} – 1.03×10^{-10}], which is around 50-fold lower than that for the mtDNA (95% HPDI: 4.8×10^{-9} – 5.4×10^{-9}). The mutation rate of the sheep MSY is slightly lower than other species: 1.4 – 4.2×10^{-10} for dog [35–38], 1.7×10^{-8} – 4.2×10^{-10} for horse [28, 39], and 1.0 – 1.9×10^{-9} for wolf [40, 41]. Bayesian skyline plot (BSP) involving 161 Y chromosome sequences showed a population increase starting ~ 0.1 mya,




Figure 2. Generation and Filtering of MSY SNPs by Using a Modified Method

(A) Multiple- and single-copy MSY contigs assembly and filtering.

(B) Confirmation of Y chromosome specificity for a contig by amplifying male (M) and female (F) genomic DNA and a no-template-control (ntc). The white bands represent the expected amplification products.

(C) Average coverage boxplot of Y chromosome genes. The dotted line below indicates a single-copy threshold, and the dotted line above indicates a multiple-copy threshold. The box plots consist of five numerical points, i.e. the minimum observed value (lower edge), 25% quantile, median, 75% quantile, and maximum observed value (upper edge). The small circles indicate outliers.

(D) The mean normalized average coverage (MNAC) of single-copy MSY (scpMSY) contigs.

(E) Average coverage of the 211 WGS individuals in scpMSY contigs and whole genome.

See also [Data S2](#) and [S3](#) and [28].

followed by a reduction after domestication (~10,000–5,000 years BP), and a more drastic decrease after 5,000 years BP (Figures 3A and 3B). During the last 2,000 years, the N_e estimates based on mitogenomes from the same 161 sheep were 47–239-fold higher than those based on the Y

chromosome, with a gradual increasing trend in the observed differences (Figures 3A and 3B).

We then analyzed an expanded sample set of 595 rams representing 118 local populations from across the world (Data S2). For the additional samples, the 81 MSY SNPs were genotyped




Figure 3. Phylogenetic Trees of 161 Domestic and 18 Wild Sheep


(A and B) Phylogenetic tree inferred from 495 MSY SNPs using Bayesian inference (BI) with posterior probabilities indicated near the nodes and Bayesian tree inferred from mitogenome sequences, respectively. See also [Figures S1B and S1C](#); [Data S4](#).

The colors at the right of the trees indicate the continental origin of the sampled individuals as in (C). Below the trees, cumulative Bayesian skyline plots are shown of MSY and mtDNA diversity based on data for 161 domestic sheep. The pink dashed lines highlight the limits of 0.28 and 1.02 mya. The short yellow and red dashed lines highlight time points of 0.01 and 0.005 mya, respectively.

(C) Neighbor-joining tree of the 179 rams based on whole-genome sequences. Colors at the right of the tree indicate the Y chromosome groups as in (A). Divergence times (million[s] years ago, mya) for the lineages were estimated on the basis of data from the 179 individuals. See also [Figure S1D](#); [Data S2](#) and [S4](#).

using the Hi-SNP genotyping platform ([Data S4](#)) and the mtDNA D-loop region (AF010406, 15,522–16,318) was sequenced (GenBank: MT768709–MT769136). We observed all paternal and maternal lineages and similar phylogenetic resolution as

above ([Figures S1E and S1F](#)). The MSY y-HA showed the highest frequency (71.1%), followed by y-HB (17.5%) and y-HC (10.8%), while y-HD was only present in four animals (0.67%) of Djallonke and Cameroon populations from Western Africa ([Data S4](#)).



(legend on next page)

Lineage y-HA is the dominant lineage in most populations (Figure 4A), while y-HB haplotypes are mainly distributed throughout semi-desertic and steppe regions in the Near East, Central Asia, Indian subcontinent, Northern Asia, and Northern Africa. Also, y-HB has a high frequency in southwest Asian and east-Mediterranean fat-tailed populations such as Barki, Chios, and Kermani (Data S5). Notably, y-HB and fat-tailed sheep have similar distribution areas [42] (Figure 4A). Out of the 39 populations carrying y-HB, 27 are fat-tailed sheep, suggesting a significant association between y-HB and the fat-tail phenotype (Pearson $\chi^2 = 7.64$, $df = 1$, $p = 0.006$) and a major influence of the colonization of fat-tailed sheep on the expansion of y-HB.

Lineage y-HC is present in Spain, England, Scandinavia, Western Russia, the Qinghai-Tibetan Plateau, and the Yunnan-Kweichow Plateau of China. The oY1 alleles in 3,056 modern samples from 214 populations reported in previous studies (Data S6) revealed a similar geographic pattern for oY1(G) (Figure 4B). Although we found oY1(G) in the two Catalanian Xisqueta and Ripollesa rams, the frequency of y-HC in these and other Spanish breeds is low (Data S6). The close similarity of the Chinese and European y-HC sequences (Figure S1B) indicate that the occurrence of y-HC in China may be partially attributed to recent introgression of English rams, which are worldwide popular breeding sires and have been used in the Yunnan Province of China [43]. In six ancient DNA samples from Finland (Medieval and post-Medieval) and Estonia (late-Chalcolithic and Medieval), only allele oY1(G) has been detected [44, 45], suggesting an ancient presence of y-HC in Northern Europe. In the same region, several sheep breeds have retained primitive features such as a small body size and short or thin tails [5], whereas a relatively high component of their genomes (up to 20%) originates from the European mouflon [46]. Thus, the range of y-HC suggests possibly an ancient expansion of primitive Northern European sheep, which became the paternal ancestors of the y-HC-carrying English mutton breeds [24], and several of them became transboundary breeds [47].

The pattern of geographic distribution may reflect ancient male founder effect and the small effective population size of males within breeds. This has led to a high frequency of y-HC in Northern Europe and of y-HB within the fat-tailed sheep. The founder effect may very well have been stimulated by selection and, for instance, have played a role in the spreading of the fat-tailed phenotype. The current evidence does not allow associating the secondary spread of wool sheep throughout the Old World with lineage y-HA or with haplotypes within this predominant lineage.

We analyzed the selected whole-genome sequences by approximate Bayesian computation (ABC) in order to obtain time estimates for the demographic expansions of the first domestic sheep and the fat-tailed sheep, which were assumed to be associated with the MSY haplogroups y-HC and y-HB, respectively. We used high-depth ($\sim 15.54\text{--}36.93 \times$ coverage) whole-genome sequences from 15 Asiatic mouflons, 16 Northern European, and 18 native fat-tailed sheep (Figure S2A; Data S2). We tested two alternative models for their expansions (Figures 4D and S2) [48]. In the best fitting model, the ancestors of Northern European primitive breeds, most of which carry y-HC, spread $\sim 10,500$ years BP (T_1 , 50% HPDI: 9,000–11,800; Figure S2F). This indicates that y-HC probably expanded as part of the dispersal of the first domestic sheep, which had a hair coat. A later spread was possibly associated with the expansion event of fat-tail sheep $\sim 3,400$ years BP (T_2 , 50% HPDI: 1,700–5,300; Figure S2F) and involved y-HB. We found that the rate of gene flow from the fat-tailed sheep to the first hair-coat domesticates ($m_2 = 2.64 \times 10^{-5}$, 50% HPDI: 5.56×10^{-6} – 1.68×10^{-4} ; Figure S2F) is higher than the rate in the opposite direction ($m_1 = 8.97 \times 10^{-6}$, 50% HPDI: 2.42×10^{-6} – 5.32×10^{-5}), which indicated a net introgression into hair sheep introducing the coarse-wool trait in most of the y-HC-carrying breeds.

These findings provided genetic support to the dispersal episodes of the first domestic hair sheep and of the fat-tailed sheep as evidenced by archeological records (Figures 1A and 1B; Data S1) and previous genetic results [5]. Although both the estimates of time under the Bayesian approximate model should be interpreted with caution because of the potential biases in the MSY mutation rate applied, the ABC results show that the fat-tailed sheep dispersed long after the spreading of the first domestic sheep and also after the emergence of the wool sheep according to the archaeological evidence (Figure 1B).

The first migratory wave of sheep populations, as inferred from the distribution of lineages y-HC and y-HD, spread to the borders of Europe and Africa as well as the highlands of Asia $\sim 10,000$ years BP. Nowadays, remnants of genetic material from the first expansion persist in Northern European sheep, which have retained a mouflon-like morphology [5], and have an Iron Age origin [49, 50] and a high proportion of their genomes originating from the European mouflon [46]. The topology of the whole-genome sequence tree (Figures 2C and S1D) suggests that these populations are related to other primitive sheep, such as the Dutch or German Heath sheep, the German Rhön, and the French Solognote and Ouessant [43], which carry y-HA. The Y chromosome of the feral European mouflons, which descend from the first domestic sheep, does not carry y-HC (Data S6). This finding

Figure 4. MSY SNP and Haplotype Frequency Distribution, Network, and Statistical Modeling

(A) Frequency distribution of the MSY haplogroups in 595 domestic sheep. Pie charts show the proportions of the haplogroups y-HA, y-HB, y-HC, and y-HD in 118 populations. The purple shade indicates the general distribution range of fat-tailed sheep [42]. See also Data S5.

(B) Distribution of Y chromosome SNP alleles oY1(G) and oY1(A) in ancient Finnish and Estonian sheep and modern sheep breeds across the world. All ancient samples show the allele oY1(G). See also Data S6.

(C) The median-joining network built with MSY data from 595 individuals. Colors indicate tail configuration (fat or thin) of the sampled sheep. See also Data S2 and S4.

(D) The best-supported demographic model for fat-tailed (FT) and primitive (PR) domestic sheep and Asiatic mouflon (AM) inferred from the approximate Bayesian computation (ABC) analysis of whole-genome sequences. N_{AM} , N_{PR} , and N_{FT} are the estimated haploid effective population sizes of Asiatic mouflons, the selected “primitive”, and fat-tailed populations of domestic sheep, respectively. T_1 and T_2 indicate the expansion time in years of sheep from the Northern European and fat-tailed populations, respectively. The numbers at the arrows indicate the gene flows between native sheep of “primitive” and fat-tailed populations. See also Figure S2; Data S2, S4, S5, and S6.

indicates that the first introduction of domestic sheep did not carry exclusively y-HC. Tibetan sheep also have primitive features but originated ~3,000 years BP [13], which is later than the earliest European sheep (Figure 1A). They carry y-HA in addition to y-HC that could have possibly been partly introduced by English rams [43]. Their whole-genome sequences cluster with those of the Asian fat-tailed sheep, suggesting gene flow between Asian breeds with different tail types. Given the multiple episodes of movement and interaction, modern populations are unlikely to act as direct proxies for the first wave of sheep that were introduced to a region. Admixture, extinction, turnover, drift, and cultural and biological processes have constantly shuffled the proportions of genetic markers in modern sheep populations in their history [51].

Our results provide genetic evidence for distinct introductions and dispersion histories for the African thin-tailed and fat-tailed populations, as previously suggested by archeological studies [52]. The fat-tailed Barki and Rahmani sheep (Egypt) and Barbary and Moboro sheep (Libya) possess the y-HB haplotypes, which are in agreement with evidence based on 50k SNP genotypes [53], indicating an immigration of Southwest Asian sheep [52]. The thin-tailed hair sheep (Mossi, Sahelian, Cameroon, and Djallonke sheep) in Western Africa can be inferred to be the first to migrate from their Near Eastern domestication center, entering the African continent via the Isthmus of Suez or across the Bab-El Mandeb strait and migrating to Western Africa overland [52]. In particular, the Djallonke and its derived population of Cameroon sheep showed an unusual y-HD haplotype, which is phylogenetically close to the haplotypes of Asiatic mouflons (Figures S1B, S1C, S1E, and S1F). Chessa et al. [5] found the same endogenous retroviral signature in the primitive European populations and the Djallonke sheep.

Overall, MSY haplotypes for the great majority of sheep populations grouped together and showed a high frequency of y-HA. Besides this, all the Merino-derived fine wool breeds carry the MSY y-HA (Data S5). Previous genetic evidence [5], together with archaeological data and historical records, suggested that the specialization for secondary products such as wool may have spurred a second expansion of sheep populations around 8,000–7,000 years BP, most likely from Southwest Asia at first [5, 54] (Figure 1B; Data S1). Thus, selection of sires from wool breeds may explain a high level of historic admixture [47, 55].

Different sheep varieties emerged over the following few thousand years. Fat-tailed sheep are known for their fat storage in their large tails and hindquarters, which provide a reserve of energy for survival in harsh production conditions with nutritional stress [56, 57]. Therefore, selection for the fat-tailed phenotype in domestic sheep and their expansion could have been initiated and accelerated by extreme climatic conditions and/or human preferences [58]. This has probably driven the third expansion in domestic sheep history from the Middle East to Northern Africa, Central and Eastern Asia, and the eastern edge of Europe, involving both y-HA and y-HB rams. The earliest archeological evidence for the history of fat-tailed sheep indicated the first description on an Uruk II stone vessel in the Near East (~5,000 years BP) [59] and the rock paintings in Ethiopia (~3,000–2,600 years BP) [60–62] (Figure 1B; Data S1). In addition, historical data record the presence of fat-tailed sheep in Central Asia since the Tsin Dynasty (266–420 AD) [63] (Data S1). In approximately the same time

period, human-mediated widespread introgression of aridity-adapted zebu (*Bos indicus*) bulls has enhanced herd survival in tropical regions [8]. The high frequencies of y-HB in modern sheep from regions of extreme environments, such as the dryland regions in Iran, Pakistan, and Afghanistan and the cold regions of Altay, Mongolia, and Siberia, coupled with archeological evidence, suggest that selection for sheep with fat tails occurred most likely first in the Near East. Thus, our data provide the first genetic evidence for a Southwest-Asian origin of fat-tailed sheep, which subsequently spread into specific regions of Asia and Africa. The observation of one y-HB haplotype (s-H22; Figure 4C) associated with the fat-tail phenotype in four European thin-tailed populations (Leccese, White Mountain, Brown Mountain, and Sumavska) may suggest introgression of fat-tailed sheep from Northern Africa via the maritime Mediterranean route [64]. In future studies, genome data should be screened for the specific traits in sheep, overcoming the limitations of neutral markers.

In conclusion, the global distribution of MSY lineages coupled with demographic reconstruction based on whole-genome sequences reveals multiple and separate waves of expansions of domestic sheep, which possibly correlate with various phenotypic traits and breeding goals for different products. At least the first domestic expansion and the dispersal of the fat-tailed sheep could have started in Southwest Asia. Our results provide important insights into the history of sheep migration and pastoralism across the Old World.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- RESOURCE AVAILABILITY
 - Lead Contact
 - Materials Availability
 - Data and Code Availability
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
- METHOD DETAILS
 - Archeological Records and Historical Data
 - Samples and Sequencing
 - Identification of MSY SNPs
 - Assembly of Mitogenomes
 - Whole-genome SNP Calling
 - MSY SNPs Genotyping
 - MtDNA D-Loop Sequencing
 - Phylogenetic Analyses
 - Geographic Patterns of the MSY Variability
 - Approximate Bayesian Computation Analysis
- QUANTIFICATION AND STATISTICAL ANALYSIS

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.cub.2020.07.077>.

ACKNOWLEDGMENTS

We are grateful to the late Zhi-Fa Wang, and Joaquim Casellas and Jordi Jordana from the Universitat Autònoma de Barcelona, and several others for

sampling. We thank Xiao-Xiao Zhang and An-Ni Yu for laboratory work and Thor Blichfeldt and the Norwegian Association of Sheep and Goat Farmers (NSG) for providing samples of Norwegian sheep. Thanks are also due to Amadou Traore (Institut de l'Environnement et de Recherches Agricoles [INERA], Ouagadougou, Burkina Faso), Masroor Ellahi Babar (Virtual University, Lahore, Pakistan), Pradeepa Silva (University of Peradeniya, Peradeniya, Sri Lanka), Seyed Abbas Rafat (University of Tabriz, College of Agriculture, Tabriz, Iran), Thiruvengadan A.K. and Saravanan Ramasamy (Tamil Nadu Veterinary and Animal Sciences University, Chennai, India) for their help in sample collection. This study was funded by grants from the National Key Research and Development Program-Key Projects of International Innovation Cooperation between Governments (2017YFE0117900), the National Natural Science Foundation of China (nos. 31825024, 31661143014, 31760661, and 31972527), the Second Tibetan Plateau Scientific Expedition and Research Program (STEP) (no. 2019QZKK0501), the External Cooperation Program of Chinese Academy of Sciences (152111KYSB20190027), the Taishan Scholars Program of Shandong Province (no. ts201511085), and Chinese Government contribution to CAAS-ILRI Joint Laboratory on Livestock and Forage Genetic Resources in Beijing and to ICARDA. The paper contributes to the CGIAR Research Program on Livestock and Fish.

AUTHOR CONTRIBUTIONS

M.-H.L. conceived the project and supervised the work; J.D. and X.-L.X. performed the majority of analysis and laboratory work with contribution from D.-F.W.; all the other co-authors provided the samples. J.D. wrote the manuscript with contributions from M.-H.L., J.-L.H., and J.A.L.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: February 20, 2020

Revised: June 14, 2020

Accepted: July 27, 2020

Published: August 20, 2020

REFERENCES

- Fages, A., Hanghoj, K., Khan, N., Gaunitz, C., Seguin-Orlando, A., Leonardi, M., McCrory Constantz, C., Gamba, C., Al-Rasheid, K.A.S., Albizuri, S., et al. (2019). Tracking five millennia of horse management with extensive ancient genome time series. *Cell* 177, 1419–1435.
- Gignoux, C.R., Henn, B.M., and Mountain, J.L. (2011). Rapid, global demographic expansions after the origins of agriculture. *Proc. Natl. Acad. Sci. USA* 108, 6044–6049.
- Brown, T.A. (1999). How ancient DNA may help in understanding the origin and spread of agriculture. *Philos. T. R. Soc. B* 354, 89–98.
- Harris, D.R., and Gosden, C. (1996). "The beginnings of agriculture in western Central Asia" in *Origins of Agriculture in Western Central Asia* (London: University College London Press).
- Chessa, B., Pereira, F., Arnaud, F., Amorim, A., Goyache, F., Mainland, I., Kao, R.R., Pemberton, J.M., Beraldi, D., Stear, M.J., et al. (2009). Revealing the history of sheep domestication using retrovirus integrations. *Science* 324, 532–536.
- Larson, G., Piperno, D.R., Allaby, R.G., Purugganan, M.D., Andersson, L., Arroyo-Kalin, M., Barton, L., Climer Vigueira, C., Denham, T., Dobney, K., et al. (2014). Current perspectives and the future of domestication studies. *Proc. Natl. Acad. Sci. USA* 111, 6139–6146.
- Colli, L., Milanese, M., Talenti, A., Bertolini, F., Chen, M., Crisà, A., Daly, K.G., Del Corvo, M., Guldbrandtsen, B., Lenstra, J.A., et al.; AdaptMap Consortium (2018). Genome-wide SNP profiling of worldwide goat populations reveals strong partitioning of diversity and highlights post-domestication migration routes. *Genet. Sel. Evol.* 50, 58–77.
- Verdugo, M.P., Mullin, V.E., Scheu, A., Mattiangeli, V., Daly, K.G., Maisano Delsar, P., Hare, A.J., Burger, J., Collins, M.J., Kehati, R., et al. (2019). Ancient cattle genomics, origins, and rapid turnover in the Fertile Crescent. *Science* 365, 173–176.
- Baird, D., Fairbairn, A., Jenkins, E., Martin, L., Middleton, C., Pearson, J., Asouti, E., Edwards, Y., Kabukcu, C., Mustafaoglu, G., et al. (2018). Agricultural origins on the Anatolian plateau. *Proc. Natl. Acad. Sci. USA* 115, E3077–E3086.
- Zeder, M.A. (2008). Domestication and early agriculture in the Mediterranean Basin: Origins, diffusion, and impact. *Proc. Natl. Acad. Sci. USA* 105, 11597–11604.
- Stiner, M.C., Buitenhuis, H., Duru, G., Kuhn, S.L., Mentzer, S.M., Munro, N.D., Pöllath, N., Quade, J., Tsartsidou, G., and Özbaşaran, M. (2014). A forager-herder trade-off, from broad-spectrum hunting to sheep management at Aşıklı Höyük, Turkey. *Proc. Natl. Acad. Sci. USA* 111, 8404–8409.
- Zhao, Y.X., Yang, J., Lv, F.H., Hu, X.J., Xie, X.L., Zhang, M., Li, W.R., Liu, M.J., Wang, Y.T., Li, J.Q., et al. (2017). Genomic reconstruction of the history of native sheep reveals the peopling patterns of nomads and the expansion of early pastoralism in east Asia. *Mol. Biol. Evol.* 34, 2380–2395.
- Hu, X.J., Yang, J., Xie, X.L., Lv, F.H., Cao, Y.H., Li, W.R., Liu, M.J., Wang, Y.T., Li, J.Q., Liu, Y.G., et al. (2019). The genome landscape of Tibetan sheep reveals adaptive introgression from argali and the history of early human settlements on the Qinghai-Tibetan plateau. *Mol. Biol. Evol.* 36, 283–303.
- Kamp, K.A., and Yoffee, N. (1980). Ethnicity in ancient Western Asia during the early second millennium BC: archaeological assessments and ethno-archaeological perspectives. *Bull. Am. Schools Orient. Res.* 237, 85–104.
- Chang, C., and Koster, H.A. (1986). Beyond bones: toward an archaeology of pastoralism. *Adv. Archaeol. Method Theory* 9, 97–148.
- Lv, F.H., Peng, W.F., Yang, J., Zhao, Y.X., Li, W.R., Liu, M.J., Ma, Y.H., Zhao, Q.J., Yang, G.L., Wang, F., et al. (2015). Mitogenomic meta-analysis identifies two phases of migration in the history of eastern Eurasian sheep. *Mol. Biol. Evol.* 32, 2515–2533.
- Demirci, S., Koban Baştanlar, E., Dağtaş, N.D., Pişkin, E., Engin, A., Ozer, F., Yüncü, E., Doğan, S.A., and Togan, I. (2013). Mitochondrial DNA diversity of modern, ancient and wild sheep (*Ovis gmelinii anatolica*) from Turkey: new insights on the evolutionary history of sheep. *PLoS ONE* 8, e81952.
- Cai, D., Tang, Z., Yu, H., Han, L., Ren, X., Zhao, X., Zhu, H., and Zhou, H. (2011). Early history of Chinese domestic sheep indicated by ancient DNA analysis of Bronze Age individuals. *J. Archaeol. Sci.* 38, 896–902.
- Decker, J.E., McKay, S.D., Rolf, M.M., Kim, J., Molina Alcalá, A., Sonstegard, T.S., Hanotte, O., Götherström, A., Seabury, C.M., Praharoni, L., et al. (2014). Worldwide patterns of ancestry, divergence, and admixture in domesticated cattle. *PLoS Genet.* 10, e1004254.
- Daly, K.G., Maisano Delsar, P., Mullin, V.E., Scheu, A., Mattiangeli, V., Teasdale, M.D., Hare, A.J., Burger, J., Verdugo, M.P., Collins, M.J., et al. (2018). Ancient goat genomes reveal mosaic domestication in the Fertile Crescent. *Science* 361, 85–88.
- Bachtrog, D. (2013). Y-chromosome evolution: emerging insights into processes of Y-chromosome degeneration. *Nat. Rev. Genet.* 14, 113–124.
- Hughes, J.F., and Page, D.C. (2015). The biology and evolution of mammalian Y chromosomes. *Annu. Rev. Genet.* 49, 507–527.
- Meadows, J.R., Hanotte, O., Drögemüller, C., Calvo, J., Godfrey, R., Coltman, D., Maddox, J.F., Marzanov, N., Kantanen, J., and Kijas, J.W. (2006). Globally dispersed Y chromosomal haplotypes in wild and domestic sheep. *Anim. Genet.* 37, 444–453.
- Meadows, J.R., Hawken, R.J., and Kijas, J.W. (2004). Nucleotide diversity on the ovine Y chromosome. *Anim. Genet.* 35, 379–385.
- Oner, Y., Calvo, J.H., and Elmaci, C. (2011). Y chromosomal characterization of Turkish native sheep breeds. *Livest. Sci.* 136, 277–280.
- Zhang, M., Peng, W.F., Yang, G.L., Lv, F.H., Liu, M.J., Li, W.R., Liu, Y.G., Li, J.Q., Wang, F., Shen, Z.Q., et al. (2014). Y chromosome haplotype diversity of domestic sheep (*Ovis aries*) in northern Eurasia. *Anim. Genet.* 45, 903–907.

27. Li, X., Yang, J., Shen, M., Xie, X.L., Liu, G.J., Xu, Y.X., Lv, F.H., Yang, H., Yang, Y.L., Liu, C.B., et al. (2020). Whole-genome resequencing of wild and domestic sheep identifies genes associated with morphological and agronomic traits. *Nat. Commun.* **11**, 2815–2830.
28. Wallner, B., Palmieri, N., Vogl, C., Rigler, D., Bozjak, E., Druml, T., Jagannathan, V., Leeb, T., Fries, R., Tetens, J., et al. (2017). Y chromosome uncovers the recent oriental origin of modern stallions. *Curr. Biol.* **27**, 2029–2035.
29. Valdez, R., Nadler, C.F., and Bunch, T.D. (1978). Evolution of wild sheep in Iran. *Evolution* **32**, 56–72.
30. Rezaei, H.R., Naderi, S., Chintauan-Marquier, I.C., Jordan, S., Taberlet, P., Virk, A.T., Naghash, H.R., Rioux, D., Kaboli, M., Luikart, G., and Pompanon, F. (2010). Evolution and taxonomy of the wild species of the genus *Ovis* (Mammalia, Artiodactyla, Bovidae). *Mol. Phylogenet. Evol.* **54**, 315–326.
31. Larson, G., and Fuller, D.Q. (2014). The evolution of animal domestication. *Annu. Rev. Ecol. Evol. Syst.* **45**, 115–136.
32. Barbato, M., Hailer, F., Orozco-terWengel, P., Kijas, J., Mereu, P., Cabras, P., Mazza, R., Pirastru, M., and Bruford, M.W. (2017). Genomic signatures of adaptive introgression from European mouflon into domestic sheep. *Sci. Rep.* **7**, 7623–7635.
33. Felkel, S., Wallner, B., Chuluunbat, B., Yadamsuren, A., Faye, B., Brem, G., Walzer, C., and Burger, P.A. (2019). A first y-chromosomal haplotype network to investigate male-driven population dynamics in domestic and wild bactrian camels. *Front. Genet.* **10**, 423–429.
34. Drummond, A.J., and Rambaut, A. (2007). BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol. Biol.* **7**, 214–221.
35. Ding, Z.L., Oskarsson, M., Ardalan, A., Angleby, H., Dahlgren, L.G., Tepeli, C., Kirkness, E., Savolainen, P., and Zhang, Y.P. (2012). Origins of domestic dog in southern East Asia is supported by analysis of Y-chromosome DNA. *Heredity* **108**, 507–514.
36. Oetjens, M.T., Martin, A., Veeramah, K.R., and Kidd, J.M. (2018). Analysis of the canid Y-chromosome phylogeny using short-read sequencing data reveals the presence of distinct haplogroups among Neolithic European dogs. *BMC Genomics* **19**, 350–358.
37. Frantz, L.A.F., Mullin, V.E., Pionnier-Capitan, M., Lebrasseur, O., Ollivier, M., Perri, A., Linderholm, A., Mattiangeli, V., Teasdale, M.D., Dimopoulos, E.A., et al. (2016). Genomic and archaeological evidence suggest a dual origin of domestic dogs. *Science* **352**, 1228–1231.
38. Botigué, L.R., Song, S., Scheu, A., Gopalan, S., Pendleton, A.L., Oetjens, M., Taravella, A.M., Seregely, T., Zeeb-Lanz, A., Arbogast, R.M., et al. (2017). Ancient European dog genomes reveal continuity since the Early Neolithic. *Nat. Commun.* **8**, 16082–16092.
39. Felkel, S., Vogl, C., Rigler, D., Dobretsberger, V., Chowdhary, B.P., Distl, O., Fries, R., Jagannathan, V., Janečka, J.E., Leeb, T., et al. (2019). The horse Y chromosome as an informative marker for tracing sire lines. *Sci. Rep.* **9**, 6095–6099.
40. Smeds, L., Kojola, I., and Ellegren, H. (2019). The evolutionary history of grey wolf Y chromosomes. *Mol. Ecol.* **28**, 2173–2191.
41. Skoglund, P., Ersmark, E., Palkopoulou, E., and Dalén, L. (2015). Ancient wolf genome reveals an early divergence of domestic dog ancestors and admixture into high-latitude breeds. *Curr. Biol.* **25**, 1515–1519.
42. Mason, I.L. (1984). *Evolution of Domesticated Animals* (London, New York: Longman).
43. Mason, I.L. (1996). *A World Dictionary of Livestock Breeds, Types and Varieties*, Fourth Edition (C.A.B International).
44. Niemi, M., Bläuer, A., Iso-Touru, T., Nyström, V., Harjula, J., Taavitsainen, J.P., Storå, J., Lidén, K., and Kantanen, J. (2013). Mitochondrial DNA and Y-chromosomal diversity in ancient populations of domestic sheep (*Ovis aries*) in Finland: comparison with contemporary sheep breeds. *Genet. Sel. Evol.* **45**, 2–15.
45. Rannamäe, E., Lõugas, L., Niemi, M., Kantanen, J., Maldre, L., Kadõrova, N., and Saarma, U. (2016). Maternal and paternal genetic diversity of ancient sheep in Estonia from the Late Bronze Age to the post-medieval period and comparison with other regions in Eurasia. *Anim. Genet.* **47**, 208–218.
46. Ciani, E., Mastrangelo, S., Da Silva, A., Marroni, F., Ferenčaković, M., Ajmone-Marsan, P., Baird, H., Barbato, M., Colli, L., Delvento, C., et al.; Econogene Consortium; Sheephapmap Consortium (2020). On the origin of European sheep as revealed by the diversity of the Balkan breeds and by optimizing population-genetic analysis tools. *Genet. Sel. Evol.* **52**, 25–38.
47. Kijas, J.W., Lenstra, J.A., Hayes, B., Boitard, S., Porto Neto, L.R., San Cristobal, M., Servin, B., McCulloch, R., Whan, V., Gietzen, K., et al.; International Sheep Genomics Consortium Members (2012). Genome-wide analysis of the world's sheep breeds reveals high levels of historic mixture and strong recent selection. *PLoS Biol.* **10**, e1001258.
48. Wegmann, D., Leuenberger, C., Neuenschwander, S., and Excoffier, L. (2010). ABCtoolbox: a versatile toolkit for approximate Bayesian computations. *BMC Bioinformatics* **11**, 116–122.
49. Long, J.L. (2004). *Introduced Mammals of the World: Their History, Distribution and Influence, Volume 79* (Wallingford (United Kingdom) and Cambridge (Massachusetts): CABI Publishing).
50. Jørgensen, G.H.M., Andersen, I.L., Holand, Ø., and Bøe, K.E. (2011). Differences in the spacing behaviour of two breeds of domestic sheep (*Ovis aries*)-influence of artificial selection? *Ethology* **117**, 597–605.
51. Pickrell, J.K., and Reich, D. (2014). Toward a new history and geography of human genes informed by ancient DNA. *Trends Genet.* **30**, 377–389.
52. Muigai, A.W., and Hanotte, O. (2013). The origin of african sheep: archaeological and genetic perspectives. *Afr. Archaeol. Rev.* **30**, 39–50.
53. Mwacharo, J.M., Kim, E.S., Elbeltagy, A.R., Aboul-Naga, A.M., Rischkowsky, B.A., and Rothschild, M.F. (2017). Genomic footprints of dryland stress adaptation in Egyptian fat-tail sheep and their divergence from East African and western Asia cohorts. *Sci. Rep.* **7**, 17647–17656.
54. Becker, C., Benecke, N., Grabundžija, A., Küchermann, H., Pollock, S., Schier, W., Schoch, C., Schrakamp, I., Schütt, B., and Schumacher, M. (2016). The Textile Revolution. Research into the Origin and Spread of Wool Production. *eTopoi, J. Ancient Studies* **6**, 102–145.
55. Kijas, J.W., Miller, J.E., Hadfield, T., McCulloch, R., Garcia-Gamez, E., Porto Neto, L.R., and Cockett, N. (2012). Tracking the emergence of a new breed using 49,034 SNP in sheep. *PLoS ONE* **7**, e41508.
56. Xu, S.S., Ren, X., Yang, G.L., Xie, X.L., Zhao, Y.X., Zhang, M., Shen, Z.Q., Ren, Y.L., Gao, L., Shen, M., et al. (2017). Genome-wide association analysis identifies the genetic basis of fat deposition in the tails of sheep (*Ovis aries*). *Anim. Genet.* **48**, 560–569.
57. Alves, S.P., Bessa, R.J.B., Quaresma, M.A.G., Kilminster, T., Scanlon, T., Oldham, C., Milton, J., Greeff, J., and Almeida, A.M. (2013). Does the fat tailed Damara ovine breed have a distinct lipid metabolism leading to a high concentration of branched chain fatty acids in tissues? *PLoS ONE* **8**, e77313.
58. Epstein, H. (1971). *The Origin of Domestic Animals of Africa* (New York, USA: Africana Publishing Corporation).
59. Ryder, M.L. (1983). *Sheep and Man* (London: Duckworth Press).
60. Finneran, N. (2007). *The Rock Art of Africa* (Routledge Press).
61. Clutton-Brock, J. (1993). *The Spread of Domestic Animals in Africa* (London: Routledge).
62. Clark, J.D., and Williams, M.A.J. (1978). Recent archaeological research in southeastern Ethiopia (1974–1975). *Annals d’Ethiopie* **11**, 19–44.
63. Feng, W.Q. (1991). A preliminary study on the formation of sheep breeds in ancient China (in Chinese). *Agric. Archaeol.* **3**, 346–353, doi: CNKI:SUN: NOSE.0.1991-03-063.
64. Pereira, F., Queirós, S., Gusmão, L., Nijman, I.J., Cuppen, E., Lenstra, J.A., Davis, S.J., Nejmeddine, F., and Amorim, A.; Econogene Consortium (2009). Tracing the history of goat pastoralism: new clues from mitochondrial and Y chromosome DNA in North Africa. *Mol. Biol. Evol.* **26**, 2765–2773.

65. Jiang, Y., Xie, M., Chen, W., Talbot, R., Maddox, J.F., Faraut, T., Wu, C., Muzny, D.M., Li, Y., Zhang, W., et al. (2014). The sheep genome illuminates biology of the rumen and lipid metabolism. *Science* **344**, 1168–1173.
66. Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120.
67. Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760.
68. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R.; 1000 Genome Project Data Processing Subgroup (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079.
69. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841–842.
70. Thorvaldsdóttir, H., Robinson, J.T., and Mesirov, J.P. (2013). Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief. Bioinform.* **14**, 178–192.
71. McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernysky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., and DePristo, M.A. (2010). The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303.
72. Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E., Lunter, G., Marth, G.T., Sherry, S.T., et al.; 1000 Genomes Project Analysis Group (2011). The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158.
73. Hahn, C., Bachmann, L., and Chevreux, B. (2013). Reconstructing mitochondrial genomes directly from genomic next-generation sequencing reads—a baiting and iterative mapping approach. *Nucleic Acids Res.* **41**, e129.
74. Kumar, S., Stecher, G., and Tamura, K. (2016). MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* **33**, 1870–1874.
75. Posada, D. (2008). jModelTest: phylogenetic model averaging. *Mol. Biol. Evol.* **25**, 1253–1256.
76. Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A., Bender, D., Maller, J., Sklar, P., de Bakker, P.I., Daly, M.J., and Sham, P.C. (2007). PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575.
77. Vilella, A.J., Severin, J., Ureta-Vidal, A., Heng, L., Durbin, R., and Birney, E. (2009). EnsemblCompara GeneTrees: Complete, duplication-aware phylogenetic trees in vertebrates. *Genome Res.* **19**, 327–335.
78. Foll, M., and Gaggiotti, O. (2008). A genome-scan method to identify selected loci appropriate for both dominant and codominant markers: a Bayesian perspective. *Genetics* **180**, 977–993.
79. Excoffier, L., and Foll, M. (2011). fastsimcoal: a continuous-time coalescent simulator of genomic diversity under arbitrarily complex evolutionary scenarios. *Bioinformatics* **27**, 1332–1334.
80. Excoffier, L., Dupanloup, I., Huerta-Sánchez, E., Sousa, V.C., and Foll, M. (2013). Robust demographic inference from genomic and SNP data. *PLoS Genet.* **9**, e1003905.
81. Excoffier, L., and Lischer, H.E. (2010). Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* **10**, 564–567.
82. Wei, T., and Simko, V. (2017). R package “corplot”: visualization of a correlation matrix (Version 0.84). <https://github.com/taiyun/corplot> 230.
83. Csillery, K., Francois, O., and Blum, M.G. (2012). abc: an R package for approximate Bayesian computation (ABC). *Methods Ecol. Evol.* **3**, 475–479.
84. Chen, K., Zhou, Y.X., Li, K., Qi, L.X., Zhang, Q.F., Wang, M.C., and Xiao, J.H. (2016). A novel three-round multiplex PCR for SNP genotyping with next generation sequencing. *Anal. Bioanal. Chem.* **408**, 4371–4377.
85. Watterson, G.A. (1975). On the number of segregating sites in genetical models without recombination. *Theor. Popul. Biol.* **7**, 256–276.
86. Beaumont, M.A., Zhang, W., and Balding, D.J. (2002). Approximate Bayesian computation in population genetics. *Genetics* **162**, 2025–2035.
87. Wegmann, D., Leuenberger, C., and Excoffier, L. (2009). Efficient approximate Bayesian computation coupled with Markov chain Monte Carlo without likelihood. *Genetics* **182**, 1207–1218.
88. Revelle, W.R. (2019). psych: procedures for psychological, psychometric, and personality research. R package version 1.9.12. (Northwestern University) <https://cran.r-project.org/web/packages/psych/index.html>.
89. Leuenberger, C., and Wegmann, D. (2010). Bayesian computation and model selection without likelihoods. *Genetics* **184**, 243–252.
90. Kass, R.E., and Raftery, A.E. (1995). Bayes Factors. *J. Am. Stat. Assoc.* **90**, 773–795.
91. Jeffreys, H. (1998). *The Theory of Probability* (Oxford: Oxford University Press).
92. Lee, M.D., and Wagenmakers, E.J. (2014). *Bayesian Cognitive Modeling: A Practical Course* (Cambridge: Cambridge University Press).
93. Gray, M.M., Wegmann, D., Haas, R.J., White, M.A., Gabriel, S.I., Searle, J.B., Cuthbert, R.J., Ryan, P.G., and Payseur, B.A. (2014). Demographic history of a recent invasion of house mice on the isolated Island of Gough. *Mol. Ecol.* **23**, 1923–1939.
94. Nater, A., Mattle-Greminger, M.P., Nurcahyo, A., Nowak, M.G., de Manuel, M., Desai, T., Groves, C., Pybus, M., Sonay, T.B., Roos, C., et al. (2017). Morphometric, behavioral, and genomic evidence for a new orangutan species. *Curr. Biol.* **27**, 3487–3498.
95. Wegmann, D., and Excoffier, L. (2010). Bayesian inference of the demographic history of chimpanzees. *Mol. Biol. Evol.* **27**, 1425–1435.

STAR★METHODS

KEY RESOURCES TABLE

| REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|--|-------------------------------------|---|
| Biological Samples | | |
| Whole blood from urial | Iran | N/A |
| Whole blood from argali | China | N/A |
| Blood samples from modern domestic sheep | Asia, Europe, Africa, South America | N/A |
| Critical Commercial Assays | | |
| Hi-SNP genotyping platform | Novogene Bioinformatics Institute | N/A |
| DEPOSITED DATA | | |
| Sheep reference genome <i>Oar v4.0</i> | [65] | https://www.ncbi.nlm.nih.gov/assembly/GCF_000298735.2 |
| Whole genome next-generation sequences (NGS) of 3 rams analyzed for raw MSY assembly | [27] | GenBank/NCBI PRJNA624020 |
| Whole genome next-generation sequences (NGS) of 31 rams and 5 ewes analyzed for scpMSY SNP calling | This paper and [27] | GenBank/NCBI PRJNA624020 and PRJNA645671 |
| Whole genome next-generation sequences (NGS) of 161 rams, 56 ewes and 25 wild sheep analyzed for whole genome analysis | This paper and [27] | GenBank/NCBI PRJNA624020 and PRJNA645671 |
| Oligonucleotides | | |
| Single-copy MSY contigs | This paper | GenBank/NCBI MT768242–MT768708 |
| Single-copy MSY SNPs | This paper | EVA/NCBI/dbSNP PRJEB39831 |
| Mitogenome assembly | This paper | GenBank/NCBI MT768063–MT768241 |
| mtDNA D-loop | This paper | GenBank/NCBI MT768709–MT769136 and MT768063–MT768241 |
| SOFTWARE AND ALGORITHMS | | |
| Trimmomatic v0.38 | [66] | http://www.usadellab.org/cms/?page=trimmomatic |
| BWA v0.7.8 | [67] | http://bio-bwa.sourceforge.net/ |
| SAMtools v0.1.19 | [68] | https://sourceforge.net/projects/samtools/files/samtools/0.1.19/ |
| CLC Genomics Workbench 11 | N/A | https://digitalinsights.qiagen.com/products-overview/analysis-and-visualization/qiagen-clc-genomics-workbench/ |
| BEDTools v2.17.0 | [69] | https://sourceforge.net/projects/bedtools/ |
| Integrative Genomics Viewer (IGV) v2.4.5 | [70] | http://software.broadinstitute.org/software/igv/ |
| R v3.5.1 | N/A | https://www.R-project.org/ |
| Genome Analysis Toolkit (GATK) v4.0 | [71] | https://software.broadinstitute.org/gatk/download/ |
| VCFtools v0.1.13 | [72] | http://vcftools.github.io/ |
| MITObim v1.9.1 | [73] | https://github.com/chrishah/MITObim |
| MIRA v4.0.2 | [73] | https://sourceforge.net/projects/mira-assembler/ |
| Picard v2.18.12 | N/A | http://broadinstitute.github.io/picard/ |
| MEGA 7.0 | [74] | https://www.megasoftware.net/ |
| jModelTest v.2.1.4 | [75] | https://code.google.com/p/jmodeltest2 |

(Continued on next page)

Continued

| REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|---------------------------|---------------------------------|---|
| BEAST v2.5.1 | [34] | https://github.com/CompEvol/beast2/issues |
| PLINK v.1.90 | [76] | http://www.cog-genomics.org/plink2/ |
| TreeBeST v.1.9.2 | [77] | http://treesoft.sourceforge.net/treebest.shtml |
| FigTree v.1.4.3 | N/A | http://tree.bio.ed.ac.uk/software/figtree/ |
| Tracer v.1.5 | N/A | http://beast.bio.ed.ac.uk/Tracer |
| ArcGIS v10.0 | ESRI, Redlands, California, USA | https://www.esri.com/zh-cn/home |
| BAYESCAN v.2.1 | [78] | http://cmpg.unibe.ch/software/BayeScan/versions.html |
| FASTSIMCOAL v.2.5.2.21 | [79, 80] | http://cmpg.unibe.ch/software/fastsimcoal/ |
| ARLUMSTAT v.3.5.2.2 | [81] | http://cmpg.unibe.ch/software/arlequin3 |
| R package <i>corrplot</i> | [82] | https://cran.r-project.org/web/packages/corrplot/corrplot.pdf |
| R package <i>abc</i> | [83] | https://cran.r-project.org/web/packages/abc/abc.pdf |
| ABCtoolbox | [48] | https://doi.org/10.1186/1471-2105-11-116 |

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Meng-Hua Li (menghua.li@cau.edu.cn).

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

The accession number for the whole genome sequencing reads of domestic and wild sheep reported in this paper is NCBI Short Reads Archive: PRJNA645671. The single-copy MSY contigs, complete mtDNA, and D-loop sequences have been deposited at GenBank under the accessions MT768242–MT768708, MT768063–MT768241, and MT768242–MT769136, respectively. SNPs identified in single-copy MSY have been submitted to NCBI/EVA/dbSNP: PRJEB39831.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

For NGS sequencing aliquots of blood samples from 164 healthy sheep (136 adult rams and 28 adult ewes) were collected as part of routine diagnostics at the sampling sites. All animal work was conducted according to a permit (no. IOZ13015) approved by the Committee for Animal Experiments of the Institute of Zoology, Chinese Academy of Sciences (CAS), China. For domestic sheep samples, animal sampling was also approved by local authorities from where the samples were taken. For the eight argali samples, sample collection was conducted under Law of the People's Republic of China on the Protection of Wildlife and relevant regulations. For urial, we collected peripheral blood sample from one captive urial after receiving authorization for research from the Department of Environmental Protection in Iran (no. 93/34089).

METHOD DETAILS

Archeological Records and Historical Data

To investigate the distribution of domestic sheep in early historical times, we compiled archeological records and historical data of domestic sheep across the world from the published literature, including the information of archeological sites, time, ancient remains and historical records associated with sheep, in particular the woolly and fat-tailed sheep (Data S1). Specifically, we performed an extensive literature review by searching for relevant publications in “National Center for Biotechnology Information,” “Web of Science” and “China National Knowledge Infrastructure” original briefings that included the keywords (i) archeological, sheep; (ii) archeological, history, wool, sheep; and (iii) archeological, history, fat tail, sheep. These queries were deemed sufficient to find the majority of relevant archeological and historical data about sheep in the world. We adopted the upper limit of the estimated

time period for archaeological sites and historical records in the plots using the ArcMap program implemented in the ArcGIS v10.0 software (ESRI, Redlands, California, USA).

Samples and Sequencing

Blood or tissue samples were collected from a total of 598 individuals comprising 561 rams and 28 ewes of domestic sheep (*O. aries*), and 9 rams of wild sheep [8 argali (*O. ammon*) and one urial (*O. vignei*)]. The domestic sheep represent 96 native populations (498 rams) and 22 improved populations (63 rams) from different geographic origins in Asia, Europe, Africa, America, and the Middle East (Data S2). The vast majority of the samples are from local populations that have been native to specific geographic areas for several hundred years and have not been admixed by commercial flocks. Genomic DNA was extracted following the standard phenol-chloroform extraction procedure or the QIAGEN's DNeasy Blood and Tissue Kit.

For whole genome sequencing, at least 0.5 µg of genomic DNA from 164 unrelated animals representing 79 populations of domestic sheep (127 rams and 28 ewes) and 2 wild species (8 argali and one urial) (NCBI: PRJNA645671; Data S2) was used to construct a library with an insert size of approximately 350 bp. Paired-end sequencing libraries were constructed according to the manufacturer's instructions (Illumina Inc., San Diego, CA, USA), and sequenced on the Illumina HiSeq X Ten sequencer (Illumina Inc.). Raw paired-end reads at a depth of 12.45–23.20 × (average depth = 18.60 ×) were sequenced (Data S2). Published whole-genome sequence data include 16 Asiatic mouflons (9 rams and 7 ewes) and 52 domestic sheep (34 rams and 18 ewes) representing 12 populations (dataset number PRJNA624020; Data S2).

Adaptors of all raw data were removed, and quality-based trimming was performed with Trimmomatic v0.38 [66] using settings “LEADING:5 || TRAILING:5 || SLIDINGWINDOW:4:15 || MINLEN:40.”

Identification of MSY SNPs

MSY de novo assembly

Reads generated through the whole-genome data of three rams from three different populations from different geographic origins (Afghanistan Waggir sample 2N of Waggir sheep, Xinjiang Duolang DLS309 sample, and Sweden sample Gotland24514; NCBI: PRJNA624020; Data S2) were mapped to the *O. aries* genome (OARv4.0, NCBI: GCA_000298735.2) [65] using BWA v0.7.8 [67]. The unmapped reads were then extracted by SAMtools v0.1.19 [68]. CLC Genomics Workbench 11 (<https://digitalinsights.qiagen.com/products-overview/analysis-and-visualization/qiagen-clc-genomics-workbench/>) was used to assemble MSY raw contigs based on the unmapped reads. After filtering the short pieces of contigs with length less than 399 bp, we obtained a total of 12,303 raw contigs (N50 = 818 bp; Figure 2A).

Identification of single-copy MSY contigs

Whole genome next-generation sequences (NGS) of 31 rams and 5 ewes (NCBI: PRJNA624020; Data S2) were mapped to the MSY_raw_contigs using BWA v0.7.8 [67]. SAMtools v0.1.19 [68] was then used to remove PCR duplicates and filter for mapping quality. The coverage percentage (covperc) per contig was calculated with BEDTools v2.17.0 [69] for each sample. Contigs that met the following criteria were retained: covperc > 0.66 in males and < 0.25 in females.

A total of 92 randomly chosen contigs were then validated by PCR using DNA from rams as a template (Figure 2B). All contigs were checked with the Integrative Genomics Viewer (IGV) v2.4.5 [70]. Afterward, 800 contigs coded by MSY_linked were retained.

To identify single-copy MSY (scpMSY) contigs, we estimated the coverage of 5 available Y chromosome genes in sheep (*DBY*, *SRY*, *UTY*, *AMELY*, and *ZFY*) [24]. The per-base coverage was computed using genomeCoverageBed in BEDTools v2.17.0 [69]. Average coverage (averagecov) for the five genes and MSY_linked in each sample genome was calculated using R v3.5.1 (<https://www.R-project.org/>). We obtain the average coverage range of 9.5–15.4 × for the four single-copy genes (*DBY*, *SRY*, *UTY*, and *AMELY*) (Figure 2C). We then normalized the averagecov of MSY contigs for each sample based on the coverage range of the 4 single-copy genes. Contigs with a mean normalized average coverage (MNAC) less than 1.5 across all the ram samples were retained, and 333 multiple-copy MSY (mcpMSY) contigs were filtered (Figure 2D). Finally, we obtained 467 single-copy MSY (scpMSY) contigs (GenBank: MT768242–MT768708). The scpMSY contigs of each sample showed that the averagecov (~7.19–17.69 ×) was near half of the coverage of whole genome sequencing (~13.58–34.11 ×) for the 31 rams (Figure 2E, Data S2).

ScpMSY SNPs calling and filtering

From a set of 211 samples, NGS reads were mapped to the MSY-linked contigs and were then realigned using the Genome Analysis Toolkit (GATK) v4.0 [71]. Variants were called in each sample separately with GATK's HaplotypeCaller using the settings “-genotyping-mode DISCOVERY||-output-mode EMIT_ALL_SITES||-min-base-quality-score 10.” Joint genotyping was performed by merging all the samples using GATK's GenotypeGvcfs. Raw variants were filtered with VCFtools v0.1.13 [72] using the settings “-minDP 2||-minQ 20||-remove-indels”. Only the SNPs called in scpMSY contigs meeting the following criteria were retained: (i) only present in at least two males and not in females; (ii) hemizygous; (iii) absence of additional SNP within 100 bp upstream and downstream of the called SNP; and (iv) passing the visual inspection by IGV v2.4.5 [70]. After filtering, a novel set of 493 scpMSY SNPs including 85 SNPs in domestic sheep and 425 in the wild species were obtained (Data S3). Further, all the 85 scpMSY SNPs in domestic sheep and 18 randomly selected scpMSY SNPs in wild species were validated by independent PCRs in three rams and three ewes.

Assembly of Mitogenomes

The clean paired-end reads were merged as one interleaved fastq file for the 161 rams of domestic sheep and 18 rams of wild sheep. The program MITObim v1.9.1 (stable-relies on MIRA v4.0.2) [73] was used to assemble the whole mitogenomes based on the

reference (GenBank: HM236174). Each assembly was repeated 3–7 times using the following code: “/MITObim.pl -start 1 -end 3 -sample file -ref OA_mt_genome -readpool file_interleaved.fastq -quick mitogenome_ref &>log.” The size of the initial mitochondrial DNA (mtDNA) assembly was around 17,000 bp. After manual curation and multiple alignments, we obtained the complete mitochondrial genome of 16,687 bp (GenBank: MT768063–MT768241).

Whole-genome SNP Calling

Clean paired-end reads of the 203 samples (179 rams and 24 ewes) (Data S2) were mapped to the sheep reference genome (Oar v4.0, NCBI: GCA_000298735.2) using the MEM module in the program BWA v0.7.8 [67] with the parameters “bwa -k 32 -M -R.” The SAMtools v0.1.19 [68] software was used to convert mapping results into the BAM format and filter the unmapped and non-unique reads. We used Picard v2.18.12 (<http://broadinstitute.github.io/picard/>) SortSam to sort the resulting bam files according to the order of chromosome coordinates, and the program Picard MarkDuplicates was used to remove duplicate reads. After the BWA alignment, a local realignment around indels was performed to correct misalignments due to the presence of indels with GATK v4.0 [71] in two steps. First, the RealignerTargetCreator command was used to determine the regions affected by local realignment. Second, IndelRealigner was used to realign the regions found above, which produced a realigned BAM file for each sample. To obtain accurate bases, Base Quality Score Recalibration (BQSR) was applied to detect systematic sequencing errors by using GATK modules, BaseRecalibrator and ApplyBQSR.

The SNP calling followed the best practice workflow recommended by GATK v4.0 [71]. In brief, SNPs were called for each sample using the GATK HaplotypeCaller module, and then a joint genotyping step for the integrated variations union which contains all samples was performed on a combined gVCF file. Filtering parameters were set as $QD < 2.0 \parallel MQ < 40.0 \parallel FS > 60.0 \parallel SOR > 3.0 \parallel HaplotypeScore > 13.0 \parallel MQRankSum < -12.5 \parallel ReadPosRankSum < -8.0 \parallel QUAL < 30$. SNPs with non-biallelic and > 40% missing calls were removed, which yielded a dataset of 31,343,261 SNPs for further analysis.

MSY SNPs Genotyping

Genotyping failed in six out from the 85 scpMSY SNPs (contig_1373_43, contig_2494_29, contig_10246_437, contig_138_365, contig_3923_760 and contig_1717_635), thus indicating that these markers were not appropriate for multiplex PCR amplification (Data S3). The remaining 79 scpMSY SNPs were genotyped in an expanded set of samples including 434 domestic sheep from 81 populations across the world (Data S2 and S4). The SNP genotyping procedure was implemented by using the Hi-SNP genotyping platform by combing three-rounds of multiplex PCR and high-throughput NGS [84]. In addition to the 79 SNPs, two previously published SNPs mapping to the *SRY* gene in domestic sheep [oY1: EU938045-Pos 88 (A/G), oY2: EU938045-Pos 460 (G/T)] [26] were also genotyped in the same set of animals.

MtDNA D-Loop Sequencing

We sequenced the mtDNA D-loop region (749 bp) of all males using a pair of primers, MSD-F 5′-ACAACACGGACTTCCCACTC-3′ (map positions 15,522–15,541 in GenBank: AF010406) and MSD-R 5′-CCAAGCATCCCCAAAATTA-3′ (map positions 16,299–16,318 in AF010406) [16]. Each PCR mixture (30 μ L total volume) contained 15 μ L 10 \times Taq Master Mix (ComWin Biotech, Beijing, China), 1 μ L DNA (30 ng/ μ L), 1 μ L of each primer (10 pmol/ μ L), and 12 μ L ddH₂O. PCR cycling conditions were 95°C for 15 min, 35 cycles at 95°C for 30 s, 60°C for 90 s, and 72°C for 30 s, and a final extension at 72°C for 7 min. PCR products were sequenced directly in both directions using the PCR primers on an ABI 3730 capillary sequencer (Applied Biosystems, Life Technologies, NY, USA). Following quality trimming and filtering, common fragments of 285 bp (15,541–15,652 and 15,960–16,132 of AF010406) of the D-loop were aligned and edited for analysis using MEGA 7.0 [74].

Phylogenetic Analyses

MSY and mitogenome haplotypes

In the 179 rams of domestic and wild sheep, 495 scpMSY SNPs (493 novel plus the two published SNPs, Data S3) were used for haplotype reconstruction. Overall, sixty-three MSY haplotypes were defined including 49 for domestic sheep (ds-H1–ds-H49), one for urial (u-H1), six for Asiatic mouflons (am-H1–am-H6), and seven for argali (a-H1–a-H7) (Data S4). In addition, 171 mtDNA haplotypes (mt-H1–mt-H171) were defined based on the 179 mitogenomes (GenBank: MT768063–MT768241).

Phylogenetics and scpMSY haplogroups

Hierarchical likelihood ratio tests were implemented to select a best-fit model of nucleotide substitution for phylogenetic analysis using jModelTest v.2.1.4 [75]. Out of 88 candidate models, the best-fit model HKY+I was chosen for the scpMSY dataset, while HKY+I+G was selected for the mitogenomes. Phylogenetic trees were inferred using the maximum likelihood (ML), maximum parsimony (MP) and neighbor-joining (NJ) methods with the HKY model in the program MEGA v.7.0 [74] and Bayesian inference (BI) in the program BEAST v2.5.1 [34]. To compare the paternal and maternal phylogenetic relationships among the rams of domestic sheep, we constructed NJ trees based on the 81 scpMSY SNPs (Data S4) and mtDNA D-loop sequences (GenBank: MT768709–MT769136 and MT768063–MT768241) using MEGA v.7.0 [74]. The bootstrap consensus trees were inferred from 100 replicates. Neighbor-joining trees based on data from the 179 rams as well as from the 49 sheep (25 rams and 24 ewes) involved in the ABC analysis were constructed based on whole-genome sequences. Single nucleotide polymorphisms with a minor allele frequency (MAF) below 0.05 and a missing call rate above 0.1 were removed (–maf 0.05 and –geno 0.1) using PLINK v.1.90 [76] from the two datasets. We then performed LD pruning by using PLINK with the commands “–indep-pairwise (50 5 0.5)” for the dataset of 179

rams and “-indep-pairwise (50 5 0.2)” for the dataset of 49 sheep. Finally, 4,262,880 SNPs and 1,689,814 SNPs segregating in 179 rams and 49 sheep, respectively, were retained for the phylogenetic reconstruction. The NJ trees were built using TreeBeST v.1.9.2 [77] with 100 bootstraps. FigTree v.1.4.3 (<http://tree.bio.ed.ac.uk/software/figtree/>) was used to visualize the NJ trees.

Empirical Bayesian estimation of T_{MRCA} and N_e

An empirical Bayesian approach that uses a prior distribution of T_{MRCA} (time to the most recent common ancestor), based on the coalescent theory, and conducts Markov chain simulations to estimate the likelihood of parameters was applied to estimate divergence times of MSY lineages. Phylogenetic reconstruction and molecular dating of lineage splitting, including estimation of T_{MRCA} for all the ovine lineages were implemented using BEAST v.2.5.1 [34]. Due to the absence of fossil records of species from the *Ovis* genus that could be used for time calibration, we used the divergence time of 2.93 million years ago (mya) between argali and domestic sheep and the time of 2.60 mya between urial and domestic sheep as *a priori* parameter, both of which were obtained from a comprehensive evolutionary analysis [16]. A strict clock was set to calculate the mutation rate. We set the Bayesian Markov chain Monte Carlo (MCMC) length to 20,000,000 steps and logging parameters every 1,000 steps. A Maximum Clade Credibility (MCC) tree was generated using a 10% burn-in with BEAST’s TreeAnnotator and drawn with FigTree v.1.4.3 (<http://tree.bio.ed.ac.uk/software/figtree/>). Convergence was confirmed by effective sampling size (ESS) > 200 using the program Tracer v.1.5 (<http://beast.bio.ed.ac.uk/Tracer>). We simulated the posterior distribution of T_{MRCA} , conditional on the two estimates of divergence time.

We performed a grid search to obtain the Watterson’s θ (θ_W) estimator for a scaled mutation rate, $\theta_W = 2N_e\mu_{bg}$, where μ_{bg} is the mutation rate per site per year [85]. R v3.2.1 (<https://www.R-project.org/>) was used to estimate θ_W ,

$$\theta_W = S / (L * \sum(1 / (n - 1))),$$

where S is the expected number of segregating sites, L is the total number of sites, and n is the number of samples. We combined all the scpMSY contigs having the called SNPs and obtained a total length of 344,382 bp. Using the above equations, we calculated the effective population size (N_e) for the lineages of scpMSY and complete mitogenomes, separately. Also, the Bayesian skyline plot (BSP) curves for the lineages of scpMSY contigs and complete mitogenomes were drawn to reconstruct the paternal and maternal historical demographies, respectively.

Geographic Patterns of the MSY Variability

First, frequencies of the four MSY lineages in the 595 rams from 118 populations were visualized using the ArcMap program implemented in the ArcGIS v10.0 software (ESRI, Redlands, California, USA) (Data S5). A total of 61 MSY haplotypes (s-H1–s-H61) assigned to the four lineages were defined by the 81 scpMSY SNPs (Data S4). The MSY SNP oY1 alleles were collected in 3,056 modern samples from 214 domestic populations and 92 wild sheep from 7 populations (4 populations of European mouflons, 2 populations of argali and 1 population of urial) in previous studies (Data S6).

Approximate Bayesian Computation Analysis

Data preparation and processing

To gain insights into the demographic history of domestic sheep with distinctive phenotypic traits (“primitive” and fat-tailed populations), we used the approximate Bayesian computation (ABC) approach to test two alternative scenarios based on the whole genome sequences. In each scenario, we considered the Asiatic mouflon (*O. orientalis*) as the wild ancestor of domestic sheep based on archeological and genetic evidence [11] and only native populations were used in this analysis in order to minimize the impact of intense and recent artificial selection. We selected individuals representative from the “primitive” short-tailed Northern European populations and from the Near Eastern fat-tailed populations. In total, 49 individuals (16 “primitive”, 18 fat-tailed, and 15 Asiatic mouflons) were included in the ABC analysis (Figure S2A, Data S2).

To match the requirements of simulations, SNPs with a minor allele frequency (MAF) lower than 0.05 and with a missing call rate higher than 0.1 were excluded (-maf 0.05 and -geno 0.1 in PLINK). To mitigate the effect of LD, we implemented LD pruning using the command “-indep-pairwise (50 5 0.5)” in the program PLINK v.1.90 [76]. Then, SNPs located 150-kb away from genes and without missing genotypes were retained, establishing an average distance of 150-kb between SNPs to minimize the effects of linkage. Since the coalescent simulations underlying ABC inference assume neutrality, we used the Bayesian approach implemented in BAYESCAN v.2.1 [78] to detect outlier SNPs with default arguments. The SNPs with a q -value lower than 0.05 as well as those mapping to the X chromosome or to the mitochondrial genome, which would exhibit reduced N_e as compared to the autosomal regions, were eliminated. After filtering, 4,438 unlinked and neutral SNPs were kept in the final dataset for the modeling analyses.

ABC inferences

Backward coalescent simulations with recombination were performed using FASTSIMCOAL v.2.5.2.21 [79, 80] under 2 models (Figure S2D). In **Model-1**, the fat-tailed sheep (N_{FT}) first diverged from the wild ancestor of domestic sheep (N_{AM}) at time T_2 , then the “primitive” sheep (N_{PR}) split from the fat-tailed sheep (N_{FT}) at time T_1 . In **Model-2**, the “primitive” sheep (N_{PR}) first diverged from the wild ancestor of domestic sheep (N_{AM}) at time T_1 , then the fat-tailed sheep (N_{FT}) split from the “primitive” sheep (N_{PR}) at time T_2 . Black arrows represent migration rates that are simulated as independent, continuous parameters.

Model parameter prior distributions are shown in Figure S2F, which were established based on previous study [13, 80]. For testing the model, we ran 5×10^5 simulations per model. A set of 24 summary statistics were computed by the program ARLSUMSTAT v.3.5.2.2 (Figure S2B) [81] for each simulation and were used to describe genetic variations within (e.g., K , mean number of alleles; H , mean heterozygosity) and between (e.g., F_{ST}) sheep populations with different phenotypic traits (Data S2). Because correlated

summary statistics can make the model simulation redundant and even produce incorrect results [86, 87], we calculated the correlations between each pair of the summary statistics using the Spearman's rho statistic by the *corr.test* function implemented in the R package *psych* [88]. The graphical representation of correlation coefficients (Spearman's rho) among the summary statistics was then obtained using a modified script of the *corrplot* function from the R package *corrplot* (Figure S2B) [82]. Ultimately, 7 informative and independent summary statistics were selected to compare the candidate models (Figures S2B and S2E).

We performed a cross-validation procedure to assess if the 7 summary statistics can provide enough statistical power to distinguish the two scenarios using the function *cv4postpr* in the R package *abc* [83]. We used 5,000 (1%) simulations closest to the observed data and randomly chose 1,000 sets of summary statistics for each model, using a single tolerance rate (tols) of 0.05 and the method 'mnlogistic' based on multinomial logistic regression. The results showed that the two models could be clearly classified in the corresponding simulations (Figure S2C).

In order to identify the best-supported model, we compared all models simultaneously using a standard ABC-GLM approach as implemented in ABCtoolbox [48] with the 5,000 (1%) simulations closest to the observed data for each model. The marginal distributions of each model were used to calculate the posterior probability of the model (PP) by using the function *postpr* in the R package *abc* with the same arguments as function *cv4postpr*, which corresponds to the proportion of the retained simulations that presented a smaller or equal likelihood under the estimated GLM as compared to the observed data [89]. We also used the marginal densities to calculate Bayes factors (BF) for each pairwise comparison between the two candidate models [48, 90–92]. The highest support was estimated for **Model-2** using both the posterior probabilities and the Bayes Factors (Figure S2D). Furthermore, we calculated the probability (*p value*) of the observed data under the general linear model (GLM) used for the post-sampling regression with ABCtoolbox [48]. A small probability indicates that the inferred GLM does not fit the observed data well. We obtained a *p value* of 1, showing that **Model-2** can reproduce the observed data well (Figure S2D) [93, 94]. As evidence, all the 7 selected observed summary statistics occurred within the 95% (i.e., 2.5%–97.5%) percentiles of the simulated summary statistics (Figure S2E).

For the demographic parameters inference, we re-ran 2×10^6 simulations under the best-supported model (**Model-2**). We retained 10,000 (0.5%) simulations that were closest to the observed data under the best-supported model (**Model-2**) (Figures 4D and S2D) following the methods described previously [48]. We then utilized the retained simulation data to calculate the median, mode, 50% and 95% highest posterior density (HPD) intervals for each demographic parameter under **Model-2** (Figure S2F). Furthermore, we checked for the biased posterior distributions based on 1,000 pseudo-observed datasets randomly chosen from the simulated data. We then computed the coverage properties of the posterior distribution using our 10,000 closest simulations. Uniformity was assessed using a classical Kolmogorov-Smirnov test using the function *ks.test* in the R package for each parameter independently (Figure S2F) [48, 87, 95]. Deviations from uniformity indicate biased posterior distributions and the corresponding parameter estimates should be considered with caution.


QUANTIFICATION AND STATISTICAL ANALYSIS

The Pearson's Chi-square test was used to assess the association between y-HB and the fat-tail phenotype. $\chi^2_{(df=1, p=0.01)} = 6.63$; $\chi^2_{(df=1, p=0.05)} = 3.84$. In the ABC analysis, Bayes factors (BF) were calculated to indicate the relative strength of evidence for two candidate models based on their own marginal densities: $1 < BF \leq 3$ (non-significant evidence); $3 < BF \leq 10$ (substantial evidence); $BF > 10$ (strong evidence). We used ABCtoolbox to compute a probability (*p value*) to compare the marginal density of the observed data with marginal densities obtained from the retained simulations, which measured the ability of the model to reproduce the observed data. $^{**}p \leq 0.01$; $^*p \leq 0.05$. Kolmogorov-Smirnov test was used to compare the posterior distributions of parameters of the best-supported model against a uniform distribution. $^{**}p \leq 0.01$; $^*p \leq 0.05$.


Supplemental Information

Paternal Origins and Migratory Episodes of Domestic Sheep


Juan Deng, Xing-Long Xie, Dong-Feng Wang, Chao Zhao, Feng-Hua Lv, Xin Li, Ji Yang, Jia-Lin Yu, Min Shen, Lei Gao, Jing-Quan Yang, Ming-Jun Liu, Wen-Rong Li, Yu-Tao Wang, Feng Wang, Jin-Quan Li, EEr Hehua, Yong-Gang Liu, Zhi-Qiang Shen, Yan-Ling Ren, Guang-Jian Liu, Ze-Hui Chen, Neena A. Gorkhali, Hossam E. Rushdi, Hosein Salehian-Dehkordi, Ali Esmailizadeh, Maryam Nosrati, Samuel R. Paiva, Alexandre R. Caetano, Ondřej Stěpánek, Ingrid Olsaker, Christina Weimann, Georg Erhardt, Ino Curik, Juha Kantanen, Joram M. Mwacharo, Olivier Hanotte, Michael W. Bruford, Elena Ciani, Kathiravan Periasamy, Marcel Amills, Johannes A. Lenstra, Jian-Lin Han, Hong-Ping Zhang, Li Li, and Meng-Hua Li




B Male-specific region of the Y chromosome



C




D



E

Male-specific region of the Y chromosome



F

mtDNA D-loop








Figure S1. Phylogenetic trees based on MSY, mitogenome and whole-genome data. Related to Figure 3.

- (A) Phylogenetic trees using Maximum likelihood (ML), maximum parsimony (MP), neighbor joining (NJ) with bootstrap values (1,000 replicates) and Bayesian inference (BI) with posterior probabilities indicated near the nodes based on the 49 MSY haplotypes, related to (see Data S4).
- (B) The phylogenetic relationship of Y-chromosomal lineages based on 495 scpMSY SNPs from 179 rams (see Data S4).
- (C) The phylogenetic relationship of mtDNA lineages based on 179 mitogenome sequences (GenBank MT768063–MT768241).
- (D) Neighbor-joining tree with 100 bootstrap replicates based on the whole-genome sequences, related to Figure 3C.
- (E) The neighbor-joining tree based on the 81 MSY SNPs from a total of 595 rams (see Data S4).
- (F) The neighbor-joining tree based on mtDNA D-loop sequences from a total of 589 rams (GenBank MT768709–MT769136 and MT768063–MT768241). Divergence times [million(s) years ago, myr] for the lineages were estimated on the basis of data from the 179 individuals.



E

| Statistic | Obs. Values | 2.5%–97.5% percentiles |
|-----------|-------------|------------------------|
| H_PR | 0.3453 | 0.0756–0.3781 |
| H_AM | 0.3678 | 0.1009–0.3988 |
| H_FT | 0.3442 | 0.0825–0.3777 |
| sd_H | 0.0133 | 0.0043–0.0325 |
| FST_AM_PR | 0.1288 | 0.0014–0.1589 |
| FST_FT_PR | 0.0597 | 0.0002–0.0833 |
| FST_FT_AM | 0.1234 | 0.0097–0.1399 |

F

| Parameter | Prior distribution | P _value_KS | Mode | Median | 50% -HPDI | 95% -HPDI |
|-----------|---------------------------|---------------|----------|----------|-----------------|-------------------|
| GAMMA | uniform (0–15) | 0.0306 | 10.1201 | 8.1195 | 6.1712–12.5375 | 1.3664–14.9550 |
| MUTATION | uniform (1e-9–1e-6) | 0.0634 | 8.89e-07 | 8.20e-07 | 3.96e-7–1.38e-6 | 8.60e-8–1.46e-6 |
| N_{AM} | uniform (1,000–1,000,000) | < 0.001 | 882,000 | 817,890 | 790,000–968,000 | 547,000–1,000,000 |
| N_{PR} | uniform (1,000–1,000,000) | 0.0127 | 27,000 | 107,203 | 1,000–107,000 | 1,000–330,000 |
| N_{FT} | uniform (1,000–1,000,000) | 0.2502 | 80,000 | 278,567 | 1,000–279,000 | 1,000–821,000 |
| m_1 | loguniform (1e-6–1e-2) | 0.1783 | 8.97e-06 | 2.60e-05 | 2.42e-6–5.32e-5 | 1.00e-6–1.80e-3 |
| m_2 | loguniform (1e-6–1e-2) | 0.3876 | 2.64e-05 | 5.05e-05 | 5.56e-6–1.68e-4 | 1.00e-6–2.91e-3 |
| T_1 | uniform (30–12,000) | < 0.001 | 10,492 | 9,234 | 9,020–11,773 | 4,293–12,000 |
| T_2 | uniform (30–12,000) | < 0.001 | 3,382 | 4,123 | 1,694–5,298 | 30–9,022 |

Figure S2. Summary information of approximate Bayesian computation (ABC) analysis for demographic inference. Related Figure 4D.

(A) The neighbor-joining tree of 34 domestic sheep and 15 Asiatic Mouflon constructed based on 1.69 million SNPs. The native sheep with different phenotypic traits and Asiatic Mouflon is highlighted using different colors. The corresponding MSY haplotypes of rams are displayed next to their samples. The individuals in gray are females.

(B) Spearman correlation plots of 24 simulated summary statistics used in the program ABCtoolbox. Ellipse glyphs are shaped to match the corresponding Spearman's coefficient. Italic and red characters represent 7 selected summary statistics, which are utilized for the model choice and parameter inference. Population specific summary statistics: the mean number of alleles over loci for single genetic group (K), the standard deviation over loci of the number of allele for single genetic group (K_{sd}), mean number of alleles over loci and genetic groups ($mean_K$), standard deviation over genetic groups of the mean number of alleles (sd_K), mean total number of alleles over loci (tot_K), mean heterozygosity over loci for single genetic group (H), standard deviation over loci of the heterozygosity for single genetic group (H_{sd}), mean heterozygosity over loci and genetic groups ($mean_H$), standard deviation of the mean heterozygosity over genetic groups (sd_H), mean total heterozygosity (tot_H). Population pairwise comparisons: pairwise F_{ST} between genetic groups (F_{ST}), mean number of differences between genetic groups (PI). PR, "Primitive" Sheep; AM, Asiatic Mouflon; FT, Fat-Tailed Sheep.

(C) Misclassification of the two demographic models using the R package abc. The confusion matrix is based on 1,000 samples ($nval$) of each model. Each grey shade from dark to light corresponds to the models 1–2.

(D) The model-testing approach compared two candidate models. In this comparison, **Model-2** is the best-fitting model. The corresponding posterior probability (PP), Bayes factor (BF) and p -value are shown above the two models, all parameters are reported in Figure S2F.

(E) Summary statistics for the 7 selected observed dataset with corresponding 2.5 and 97.5 percentiles. Distributions are generated from the 5,000 (1%) simulations closest to the observed dataset of **Model-2**.

(F) Parameter estimation of the best supported model (**Model-2**) in the approximate Bayesian computation (ABC) analysis. GAMMA, gamma distribution of the mutation rate; MUTATION, rate per site per generation, with a generation time assumed to be 3 years; N_{AM} , the estimated effective population size of Asiatic Mouflon, in haploids; N_{PR} , the estimated effective population size of "Primitive" sheep, in haploids; N_{FT} , the estimated effective population size of Fat-Tailed Sheep, in haploids; m_1 , migration rate from "Primitive" sheep (N_{PR}) to Fat-Tailed sheep (N_{FT}); m_2 , migration rate from Fat-Tailed Sheep (N_{FT}) to "Primitive" Sheep (N_{PR}); T_1 , the split time of "Primitive" sheep, in years; T_2 , the split time of Fat-Tailed sheep, in years; P_value_KS , the P value of the Kolomogorov-Smirnov test of uniformity of the posterior quantiles; 50%-HPD, 50%-highest posterior density interval; 95%-HPD, 95%-highest posterior density interval. All other models use the same prior bound.

Supplemental References

- S1. Magnavita, S. (2008). The Oldest Textiles from Sub-Saharan West Africa: Woolen Facts from Kissi, Burkina Faso. *J. Afr. Archaeol.* 6(2), 243–254.
- S2. Gardiner, A. (1961). *Egypt of the pharaohs*, (Cambridge: Cambridge University Press).
- S3. Pleurdeau, D., Imalwa, E., Detroit, F., Lesur, J., Veldman, A., Bahain, J.J., and Marais, E. (2012). "Of sheep and men": earliest direct evidence of caprine domestication in southern Africa at Leopard Cave (Erongo, Namibia). *PloS one* 7, e40340.
- S4. Finneran, N. (2007). *The rock art of Africa*. (Routledge Press).
- S5. Dunne, J., Evershed, R.P., Salque, M., Cramp, L., Bruni, S., Ryan, K., Biagetti, S., and di Lernia, S. (2012). First dairying in green Saharan Africa in the fifth millennium BC. *Nature* 486, 390–394.
- S6. Clutton-Brock, J. (1993). The spread of domestic animals in Africa. In T. Shaw, P. Sinclair, B. Andah, and A. Okpoko, eds. (London: Routledge).
- S7. Clutton-Brock, J. (1999). *A natural history of domesticated mammals*. (Cambridge: Cambridge University Press).
- S8. Clark, J.D., and Williams, M.A.J. (1978). Recent archaeological research in southeastern Ethiopia (1974–1975). *Annals d’Ethiopie* 11, 19–44.
- S9. Greco, E., El-Aguizy, O., Ali, M.F., Foti, S., Cunsolo, V., Saletti, R., and Ciliberto, E. (2018). Proteomic analyses on an ancient Egyptian cheese and biomolecular evidence of brucellosis. *Anal. Chem.* 90, 9673–9676.
- S10. Gautier, A. (1987). Fishing, fowling and hunting in Late Palaeolithic times in the Nile Valley in Upper Egypt. *Palaeoecol. Afr.* 18, 429–440.
- S11. Close, A.E. (1992). Holocene occupation of the Eastern Sahara. In Klees, F., and Kuper, R., eds. (Heinrich Barth Institute, Koln: New Light on the Northeast African Past).
- S12. Vermeersch, P.M., Van Peer, P., Moeyersons, J., and Van Neer, W. (1996). *Neolithic occupation of the Sodmein area, Red Sea Mountains, Egypt*. (Harare: University of Zimbabwe Publications).
- S13. Oren, E.D. (1979). Die Landbrücke zwischen Asien und Afrika. *Archäologie der Nordsinai bis zur klassischen Periode*. Rothenberg, B.I., ed. (Bern: Kümmerly und Frey).
- S14. Chaix, L., and Grant, A. A study of a pre-historic population of sheep (*Ovis aries*) from Kerma. *Archaeozoologia* 1, 93–107.
- S15. Iwaszczuk, U., Niderla-Bielinska, J., and Scieczynska, A. (2019). Kings and peasants from El-Zuma/El-Detti microregion in the Early Makurian period. Economic aspects of animal bones from funerary contexts. *PloS one* 14, e0212423.
- S16. Barthelme, J.W. (1985). Fisher-hunters and Neolithic pastoralists in East Turkana, Kenya. (Oxford: British Archaeological Reports, International Series 254).
- S17. MacDonald, K.C., and MacDonald, R.H. (2000). *The origins and development of domesticated animals in arid West Africa*. (New York: Routledge).

- S18. Vermeersch, P.M., Van Peer, P., Moeyersons, J., and Van Neer, W. (1994). Sodmein Cave site, Red Mountains. *Sahara* 6, 32–40.
- S19. Henshilwood, C. (1996). A revised chronology for pastoralism in southernmost Africa: New evidence of sheep at c. 2000 b.p. from Blombos Cave, South Africa. *Antiquity*, 945–949.
- S20. Rocha, J., Chen, S., and Beja-Pereira, A. (2011). Molecular evidence for fat-tailed sheep domestication. *Trop. Anim. Health. Pro.* 43, 1237–1243.
- S21. Soe, M.J., Nejsun, P., Seersholm, F.V., Fredensborg, B.L., Habraken, R., Haase, K., Hald, M.M., Simonsen, R., Hojlund, F., Blanke, L., et al. (2018). Ancient DNA from latrines in Northern Europe and the Middle East (500 BC–1700 AD) reveals past parasites and diet. *PloS one* 13, e0195481.
- S22. Itahashi, Y., Erdal, Y.S., Tekin, H., Omar, L., Miyake, Y., Chikaraishi, Y., Ohkouchi, N., and Yoneda, M. (2019). Amino acid ¹⁵N analysis reveals change in the importance of freshwater resources between the hunter-gatherer and farmer in the Neolithic upper Tigris. *Am. J. Phys. Anthropol.*, 1–11.
- S23. Anne, W.T., Muigai, and Olivier, H. (2013). The origin of African sheep: archaeological and genetic perspectives. *Afr. Archaeol. Rev.* 30, 39–50.
- S24. Good., I.L. (1999). *The Ecology of Exchange: Textiles from Shahr-i Sokhta, Eastern Iran.* (ProQuest Dissertations and Theses).
- S25. Shishlina, N.I., Orfinskaya, O.V., and Golikov, V.P. (2003). Bronze Age Textiles from the North Caucasus: New Evidence of Fourth Millennium BC Fibres and Fabrics. *Oxford J. Archaeol.* 22 (4), 331–344.
- S26. Baird, D., Fairbairn, A., Jenkins, E., Martin, L., Middleton, C., Pearson, J., Asouti, E., Edwards, Y., Kabukcu, C., Mustafaoglu, G., et al. (2018). Agricultural origins on the Anatolian plateau. *P. Natl. Acad. Sci. U.S.A.* 115, E3077–E3086.
- S27. Bokonyi, S. (1977a). The Animal Remains of Four Sites in the Kermanshah Valley, Iran: Asiab, Dehsavar and Siahbid: The Faunal Evolution, Environmental Changes and Development of Animal Husbandry VII–III Millennia BC. Oxford: Brit. Archaeol. Rep. 34.
- S28. Bokonyi, S. and Bartosiewicz, L. (1997b). Tierknochenfunde. In *Karanovo-Die Ausgrabungen im Südsektor 1984–1992*. Ed. by S. Hiller and V. Nikolov. Österreichisch Bulgarische Ausgrabungen und Forschungen in Karanovo I.1. Horn/Wien: Verlag Ferdinand Berger & Söhne, 385–480.
- S29. Benecke, N. Tierhaltung und Jagd in den mittel-und spätneolithischen Siedlungen von Aşağı Pınar. (Unpublished).
- S30. Becker, C., Benecke, N., Grabundžija, A., Küchelmann, H., Pollock, S., Schier, W., Schoch, C., Schrakamp, I., Schütt, B., and Schumacher, M. (2016). The Textile Revolution. Research into the Origin and Spread of Wool Production. *eTopoi, J. Ancient Studies* 6, 102–145.
- S31. Stiner, M.C., Buitenhuis, H., Duru, G., Kuhn, S.L., Mentzer, S.M., Munro, N.D., Pollath, N., Quade, J., Tsartsidou, G., and Ozbasaran, M. (2014). A forager-herder trade-off, from broad-spectrum hunting to sheep management at Asikli Hoyuk, Turkey. *P. Natl. Acad. Sci. U.S.A.* 111, 8404–8409.
- S32. Benecke, N. (2012). Haustierhaltung und Jagd. In *Die frühbronzezeitliche Siedlung von Kanlıgeçit bei Kırklareli*. Ed. by M. Özdoğan and H. Parzinger. Archäologie in Eurasien 27, Mainz: Zabern, 249–266.

- S33. Demirci, S., Koban Bastanlar, E., Dagtas, N.D., Piskin, E., Engin, A., Ozer, F., Yuncu, E., Dogan, S.A., and Togan, I. (2013). Mitochondrial DNA diversity of modern, ancient and wild sheep (*Ovis gmelinii anatolica*) from Turkey: new insights on the evolutionary history of sheep. *PloS one* 8, e81952.
- S34. Dymova, M.A., Zadorozhny, A.V., Mishukova, O.V., Khrapov, E.A., Druzhkova, A.S., Trifonov, V.A., Kichigin, I.G., Tishkin, A.A., Grushin, S.P., and Filipenko, M.L. (2017). Mitochondrial DNA analysis of ancient sheep from Altai. *Anim. Genet.* 48, 615–618.
- S35. Gholamreza, M., Kobra, M., Mahsa, S.M., Iraj, M., Mohammad, M., Reza, N., Ghasem, H., Parisa, N., and Santiago, M.–C. (2015). *Dicrocoelium dendriticum* found in a Bronze Age cemetery in western Iran in the pre-Persepolis period: The oldest Asian palaeofinding in the present human infection hottest spot region. *Parasitol. Int.* 64, 251–255.
- S36. Taylor, W., Shnaider, S., Abdykanova, A., Fages, A., Welker, F., Irmer, F., Seguin-Orlando, A., Khan, N., Douka, K., Kolobova, K., et al. (2018). Early pastoral economies along the Ancient Silk Road: Biomolecular evidence from the Alay Valley, Kyrgyzstan. *PloS one* 13, e0205646.
- S37. Hermes, T.R., Frachetti, M.D., Doumani Dupuy, P.N., Mar'yashev, A., Nebel, A., and Makarewicz, C.A. (2019). Early integration of pastoralism and millet cultivation in Bronze Age Eurasia. *Proceedings. Biol. Sci.* 286, 20191273.
- S38. Vigne, J.D., Helmer, D., and Peters, J. (2005). *The First Steps of Animal Domestication*. (Vigne JD, Peters J, Helmer D, Ed), (Oxford: Oxbow Books).
- S39. Arnold, E.R., Hartman, G., Greenfield, H.J., Shai, I., Babcock, L.E., and Maeir, A.M. (2016). Isotopic Evidence for Early Trade in Animals between Old Kingdom Egypt and Canaan. *PloS one* 11, e0157650.
- S40. Jeong, C., Wilkin, S., Amgalantugs, T., Bouwman, A.S., Taylor, W.T.T., Hagan, R.W., Bromage, S., Tsolmon, S., Trachsel, C., Grossmann, J., et al. (2018). Bronze Age population dynamics and the rise of dairy pastoralism on the eastern Eurasian steppe. *P. Natl. Acad. Sci. U.S.A.* 115, E11248–E11255.
- S41. Atici, L., Pilaar Birch, S.E., and Erdogu, B. (2017). Spread of domestic animals across Neolithic western Anatolia: New zooarchaeological evidence from Ugurlu Hoyuk, the island of Gokceada, Turkey. *PloS one* 12, e0186519.
- S42. Losch, S., Grupe, G., and Peters, J. (2006). Stable isotopes and dietary adaptations in humans and animals at pre-pottery Neolithic Nevalı Cori, southeast Anatolia. *Am. J. Phys. Anthropol.* 131, 181–193.
- S43. Ma, Y., Fuller, B.T., Wei, D., Shi, L., Zhang, X., Hu, Y., and Richards, M.P. (2016). Isotopic perspectives ($\delta(13)C$, $\delta(15)N$, $\delta(34)S$) of diet, social complexity, and animal husbandry during the proto-shang period (ca. 2000–1600 BC) of China. *Am. J. Phys. Anthropol.* 160, 433–445.
- S44. Flad, R.K., Yuan, J., and Li, S. 2007. Zooarchaeological evidence for animal domestication in northwest China. *Dev. Quat. Sci.* 9, 167–203.
- S45. He, S.Q. (1987). A summary of the Pre-Qin agricultural archaeology in Gansu Province. *Agr. Archaeol.* 1, 55–61.
- S46. Xie, D.J. (1980a). The excavation of Zhangjiazui and Jijiachuan sites, Yongjing, Gansu Province. *Acta. Archaeologica. Sinica.* 2, 187–220.

- S47. Xie, D.J. (1980b). The Lianhuatai site of Xindian culture, Yongjing, Gansu Province. *Archaeol.* 4, 296–310.
- S48. Bo, W.C. (1987). The origin of Tibetan sheep (in Chinese). *Agr. Archaeol.* (1), 388–393.
- S49. Liu, L. and Chen, X. (2012). *The Archaeology of China: From the Late Paleolithic to the Early Bronze Age.* (Cambridge: Cambridge University Press).
- S50. Li, Z.P., Brunson, K., and Dai, L.L. (2014) The zooarchaeological study of wool exploitation from the Neolithic Age to the early Bronze Age in the central plains (in Chinese). *Quat. Sci.* 34(1), 149–156.
- S51. Chen, X.L., Fang, Y.M., Hu, Y.W., Hou, Y.F., Lü, P., Yuan, J., Song, G.D., Fuller, B.T., and Richards, M.P. (2016). Isotopic reconstruction of the late Longshan period (ca. 4200–3900 BP) dietary complexity before the onset of state-level societies at the Wadian site in the Ying River Valley, Central Plains, China. *Int. J. Osteoarchaeol.* 26, 808–817.
- S52. Dai, L.L., Li, Z.P., Zhao, C.Q., Yuan, J., Hou, L.L., Wang, C.S., Fuller, B.T., and Hu, Y.W. (2016). An isotopic perspective on animal husbandry at the Xinzhai site during the initial stage of the Legendary Xia Dynasty (2070–1600 BC). *Int. J. Osteoarchaeol.* 26, 885–896.
- S53. Dodson, J., Dodson, E., Banati, R., Li, X., Atahan, P., Hu, S., Middleton, R.J., Zhou, X., and Nan, S. (2014). Oldest directly dated remains of sheep in China. *Sci. Rep.* 4, 7170–7173.
- S54. Cai, D.W., Tang, Z.W., Yu, H.X., Han, L., Ren, X.Y., Zhao, X.B., Zhu, H., and Zhou, H. (2011). Early history of Chinese domestic sheep indicated by ancient DNA analysis of Bronze Age individuals. *J. Archaeol. Sci.* 38, 896–902.
- S55. Chen, F.H., Dong, G.H., Zhang, D.J., Liu, X.Y., Jia, X., An, C.B., Ma, M.M., Xie, Y.W., Barton, L., Ren, X.Y., et al. (2015). Agriculture facilitated permanent human occupation of the Tibetan Plateau after 3600 B.P. *Science* 347, 248–250.
- S56. Yuan, J. (2010). Zooarchaeological study on the domestic animals in ancient China. *Quat. Sci.* 30, 298–306.
- S57. Chen, X.L., Yuan, J., Hu, Y.W., He, N., and Wang, C.S. (2012). Domestic animal raising strategies 588 at the Taosi site from the isotopic perspective. *Archaeology* 540, 75–82.
- S58. Brunson, K., He, N., Dai, X. (2016). Sheep, cattle, and specialization: new zooarchaeological perspectives on the Taosi Longshan. *Int J Osteoarchaeol.* 26, 460–475.
- S59. Aldenderfer, M. (2011). Peopling the Tibetan Plateau: insights from archaeology. *High. Alt. Med. Biol.* 12, 141–147.
- S60. Fu, D.X. (2001). The discovery, identification and research of Neolithic crop remains from Changguogou site, Tibet (in Chinese). *Archaeol.* 3, 66–74.
- S61. Aldenderfer, M., Zhang, Y. (2004). The prehistory of the Tibetan Plateau to the seventh century A.D.: perspectives and research from China and the West since 1950. *J. World Prehist.* 18, 1–55.
- S62. Wagner, M., Wu, X., Tarasov, P., Aisha, A., Ramsey, C.B., Schultz, M., Schmidt-Schultz, T., and Gresky, J. (2011). Radiocarbon-dated archaeological record of early first millennium B.C. mounted pastoralists in the Kunlun Mountains, China. *P. Natl. Acad. Sci. U.S.A.* 108, 15733–15738.

- S63. You, Y., Lv, P., Wang, J.X., Ma, J., and Ren, M. (2016) The emergence and early utilization of domestic sheep in xinjiang (in Chinese). *Chinese Archaeol.* (12), 104–114.
- S64. Wang, B.H. (1983). Excavation and preliminary study of ancient tomb ditch in kongquehe river (in Chinese). *Social Sciences in Xinjiang* (1), 117–126.
- S65. Hong, C., Jiang, H., Lu, E., Wu, Y., Guo, L., Xie, Y., Wang, C., and Yang, Y. (2012). Identification of milk component in ancient food residue by proteomics. *PloS one* 7, e37053.
- S66. Yidilisi, Li, W.Y., and Hu, X.J. (2007). Xinjiang lop nur river cemetery excavation in 2003 (in Chinese). *Cult. Relics* (10), 2–3 and 6–44.
- S67. O'Sullivan, N.J., Teasdale, M.D., Mattiangeli, V., Maixner, F., Pinhasi, R., Bradley, D.G., and Zink, A. (2016). A whole mitochondria analysis of the Tyrolean Iceman's leather provides insights into the animal sources of Copper Age clothing. *Sci. Rep.* 6, 31279–31287.
- S68. Gleba, M. (2012). From textiles to sheep: investigating wool fibre development in pre-Roman Italy using scanning electron microscopy (SEM). *Journal of Archaeol. Sci.* 39(12), 3643–3661.
- S69. Ethier, J., Banffy, E., Vukovic, J., Leshtakov, K., Bacvarov, K., Roffet-Salque, M., Evershed, R.P., and Ivanova, M. (2017). Earliest expansion of animal husbandry beyond the Mediterranean zone in the sixth millennium BC. *Sci. Rep.* 7, 7146–7156.
- S70. Georgiev, G.I. (1979). Ezero. (Eine Siedlung aus der Frühbronzezeit. Sofa: Bulgarische Akademie).
- S71. Greenfield, H.J. (1988). The origins of milk and wool production in the Old World: A zooarchaeological perspective from the Central Balkans. *Curr. Anthropol.* 29(4), 573–593.
- S72. Frei, K.M., Mannering, U., Berghe, I.V., and Kristiansen, K. (2017). Bronze Age wool provenance and dye investigations of Danish textiles. *Antiquity* 91(357), 640–654.
- S73. Brandt, L.O., Schmidt, A.L., Mannering, U., Sarret, M., Kelstrup, C.D., Olsen, J.V., and Cappellini, E. (2014). Species identification of archaeological skin objects from Danish bogs: comparison between mass spectrometry-based peptide sequencing and microscopy-based methods. *PloS one* 9, e106875.
- S74. Randsborg, K. and Christensen, K. (2006). Bronze Age oak-coffin graves. *Archaeology and dendro-dating. Acta. Archaeol.* 77, 1–246.
- S75. Tams, K.W., Jensen Soe, M., Merkyte, I., Valeur Seersholm, F., Henriksen, P.S., Klingenberg, S., Willerslev, E., Kjaer, K.H., Hansen, A.J., and Kapel, C.M.O. (2018). Parasitic infections and resource economy of Danish Iron Age settlement through ancient DNA sequencing. *PloS one* 13, e0197399.
- S76. Lang, V. (2007). The Bronze and Early Iron Ages in Estonia. (*Estonian Archaeology* 3, Tartu: Tartu University Press).
- S77. Mandel, M. (2007). Von den Forschungen am Bodendenkmal zu Uugla. *Archaeological Fieldwork in Estonia* 2006, 113–119.
- S78. Russow, E. and Haak, A. (2013a). Archaeological fieldwork in 2012. *Archaeological Fieldwork in Estonia* 2012, 9–28.
- S79. Lõugas, L., Maldre, L., Tomek, T., and Kadakas, V. (2011). Archaeozoological evidence from the Padise

monastery. *Archaeological Fieldwork in Estonia* 2012, 83–92.

- S80. Lõugas, L. (1992). Pärnu Malmö tänav 15 arheoloogilised päästekaevamised 1992. (Report on the archaeozoological analysis, manuscript in the archive of the Institute of History, Tallinn University).
- S81. Kadakas, V., Haak, A., Russow, E., Saluäär, U., and Sarv, K. (2003). Archaeological Investigations in Pärnu. *Archaeological Fieldwork in Estonia* 2002, 179–209.
- S82. Tõnisson, E. (2008). Eesti muinaslinnad. Muinasaja teadus 20. (Tartu-Tallinn: Tartu Ülikooli Kirjastus).
- S83. Tõnisson, E. and Selirand, J. (1978). = Тыниссон Э., Селиранд Ю. О раскопках городища Варбола. Eesti NSV Teaduste Akadeemia Toimetised. Ühiskonnateadused 27(4), 358–360.
- S84. Kadakas, V., Nurk, R., Püüa, G., Toos, G., Lõugas, L., Hiie, S., and Kihno, K. (2010). Rescue excavations in Tallinn Vabaduse Square and Ingermanland bastion 2008–2009. *Archaeological Fieldwork in Estonia* 2009, 49–72.
- S85. Peets, J. and Maldre, L. (1995). Eesti kohaliku lambatõu kujunemisest arheoloogilise ja osteoloogilise materjali põhjal ehk neljasarvelised lambad ning Jakobsoni must kuub. *Kleio. Ajaloo ajakiri* 1(11), 3–4.
- S86. Sokolovski, V. (1997). Aruanne arheoloogilistest uuringutest Tallinnas, Roosikrantsi tn 9 ja 11, I. Uuringute tulemused. (Excavation report, manuscript in the archaeology archive of the Tallinn University).
- S87. Tiko, A. (2014). Keskaegsed majapidamised Tallinnas Sauna tn 8/10 kaevamiste näitel. (MA Thesis, Tallinn University).
- S88. Russow, E., Lõugas, L., Maldre, L., Hiie, S., Kihno, K., Luik, H., and Kadakas, V. (2013b). Medieval and early modern suburban site in Tallinn, Tartu Rd. 1: artefacts and ecofacts. *Archaeological Fieldwork in Estonia* 2012, 149–170.
- S89. Lang, V., Tvauri, A., and Rohtla, M. (2002). The hill-fort of Keava. *Arheoloogilised välitööd Eestis=Archaeological Fieldwork in Estonia* 2001, 65–73.
- S90. Laneman, M., Lang, V., Malve, M., Rannamäe, E. (2015). New data on Jaani stone graves at Vão, northern Estonia. *Estonian J. Archaeol.* 19(2), 110–137.
- S91. Konsa, M. and Kivi, K. (2012). Village at the foot of the fort: settlement sites I–III at Linnaaluste. Keava–'The Hand of the Sun'. *Estonian J. Archaeol. Supplementary series* 1, 63–91.
- S92. Tvauri, A. (2011a). Aruanne Paide Vallimäel Kõlakoja rekonstrueerimise ja mänguväljaku rajamise juures 2001. aastal toimunud arheoloogilisest järeelvalvest. (Excavation report, manuscript in the archaeology archive of the University of Tartu).
- S93. Lavi, A. (1978). Aruanne uurimistöödest Olustvere asulakohal 5.09.-15.10.1978. (Excavation report, manuscript in the archaeology archive of the Tallinn University).
- S94. Rannamae, E., Lõugas, L., Niemi, M., Kantanen, J., Maldre, L., Kadorova, N., and Saarma, U. (2016). Maternal and paternal genetic diversity of ancient sheep in Estonia from the Late Bronze Age to the post-medieval period and comparison with other regions in Eurasia. *Anim. Genet.* 47, 208–218.
- S95. Valk, H. (1999a). Aruanne arheoloogilistest uurimistöödest Viljandis Suusahüppemäe piirkonnas 04.08.-06.09. (An excavation report. Manuscript in the archaeology archive of University of Tartu).

- S96. Valk, H. (1991) Aruanne arheoloogilistest kaevamistest Viljandi Jaani kiriku kommunikatsioonidetrassil. (An excavation report. Manuscript in the archaeology archive of the University of Tartu).
- S97. Valk, H. (1993). Aruanne arheoloogilistest kaevamistest Viljandi Noorte Huvikeskuse küttetrassil 1991. a. (An excavation report. Manuscript in the archaeology archive of University of Tartu).
- S98. Haak, A. and Rannamäe, E. (2014). Tracing the castle crew. (Zoo)archaeological search for the inhabitants of Viljandi castle (South Estonia) in the late 13th century. In: *Castrum Bene* 12, Predovnik, K., ed. (University Press, Faculty of Arts, Ljubljana), pp. 139–152.
- S99. Haak, A. (2004). Archaeological investigations at Viljandi castle of the Teutonic Order and in Medieval Viljandi. *Arheoloogilised välitööd Eestis = Archeological Fieldwork in Estonia* 2003, 107–122.
- S100. Tvauri, A. (1999a). Aruanne arheoloogilistest päästekaevamistest Viljandi Spordihoone juurdeehituse alal 1999. aastal. (An excavation report. Manuscript in the archaeology archive of University of Tartu).
- S101. Haak, A. and Valk, H. (2002a). Archaeological investigations of medieval and post-medieval Viljandi. *Arheoloogilised välitööd Eestis = Archaeological Fieldwork in Estonia* 2001, 91–104.
- S102. Tvauri, A. (2008). Archaeological investigations at the Iron Age and Medieval settlement site of Sargvere. *Archaeological*, 109–114.
- S103. Tvauri, A. (1999b). Archaeological excavations in the gate building of Põltsamaa castle. *Archaeological Fieldwork in Estonia* 1998, 127–131.
- S104. Kaldre, H., Aguraiuja, Ü., and Livin, L. (2010). Preliminary investigations of fossil field systems at Loo. *Archaeological Fieldwork in Estonia* 2009, 73–84.
- S105. Mäesalu A. (1993). Die Burg Otepää als ein Zentrum in Südostestland im 11.-14. Jh. In: Drake K, editor. *Castella maris Baltici I*. (Stockholm: Almqvist & Wiksell International), pp.143–148.
- S106. Jaanits, L., Laul, S., Lõugas, V., and Tõnisson, E. (1982). Eesti esiajalugu. (Tallinn: Eesti Raamat).
- S107. Valk, H. (1989a). Magistraalkraavist väljatõstetud mullavallide sorteerimisest Otepää linnamäe ja kiriku vahel 1989. a. (Excavation report, manuscript in the archaeology archive of the University of Tartu).
- S108. Tamla T. (1984). Первое городище и хронология комплекса археологических памятников в Пада. *Eesti NSV Teaduste Akadeemia Toimetised. Ühiskonnateadused* 33(4), 360–363.
- S109. Tamla, T. (1983). Селище в Пада. *Eesti NSV Teaduste Akadeemia Toimetised. Ühiskonnateadused* 32(4), 302–306.
- S110. Kriiska, A., Roog, R., Alttoa, K., Allik, A., Läänelaid, A., Bernotas, R., and Malve, M. (2011). The medieval and modern period building complex at 2 Lutsu Street in Tartu. Results of the archaeological, architectural historical, dendrochronological and osteoarchaeological research. *Ajalooline Ajakiri* 1(135), 3–40.
- S111. Lillak, A. and Valk, H. (2009a). Archaeological trial excavations on Rõuge hill fort, South-East Estonia. *Archaeological Fieldwork in Estonia* 2008, 72–81.
- S112. Valk H. (2006). Excavations in the Late Iron Age and medieval Centres of Võrumaa: Tilleoru, Kirumpää and Vastseliina. *Archeological Fieldwork in Estonia* 2005, 127–140.
- S113. Valk, H. (2007). Excavations in the ruins of Vastseliina Castle and on the hillforts of Urvaste and Hinniala.

Archeological Fieldwork in Estonia 2001, 49–67.

- S114. Kriiska, A. and Lõhmus, M. (2006). Archaeological excavations on Suur street, Narva Town. *Archaeological Fieldwork in Estonia 2005*, 189–206.
- S115. Rammo, R. and Veldi, M. (2005). Archaeological Excavations at Musumägi Hill in Viljandi. *Archaeological Fieldwork in Estonia 2004*, 103–116.
- S116. Haak, A., Rannamäe, E. (2012). A Household by the Market Place: Archaeological and Zooarchaeological Contributions to the Development of Urban Viljandi, Estonia. In: Viitanen, E.M., ed. (18th annual meeting, European Association of Archaeologists, 29 August-1 September 2012 Helsinki, Finland, abstracts. Helsinki: University of Helsinki), p94.
- S117. Tvauri, A. (1998). Archaeological investigations in the old part of Viljandi. *Archaeological Fieldwork in Estonia 1997*, 81–86.
- S118. Haak, A. (2002b). Aruanne arheoloogilistest päästekaevamistest Viljandis, AS Domotex keskküttetrassi alal Posti tn 16 ja Lossi tn 33 vahelises hoovis 31. oktoobrist 17. Novembrini 2001. a. (An excavation report. Manuscript in the archaeology archive of University of Tartu).
- S119. Valk, H., Pluskowski, A., Brown, A.D., Rannamäe, E., Malve, M., and Varul, L. (2011a). The castle of the Teutonic Order in Karksi: preliminary excavation results. *Archaeological Fieldwork in Estonia*, 47–56.
- S120. Valk, H., Rannamäe, E., Brown, A.D., Pluskowski, A., Badura, and M., Lõugas, L. (2012). Thirteenth century cultural deposits at the castle of the Teutonic Order in Karksi. *Archaeological Fieldwork in Estonia*, 73–92.
- S121. Valk, H., Kama, P., Olli, M., and Rannamae, E. (2011b). Excavations on the hill forts of south–eastern Estonia: Kõivuküla, Märdi, Truuta and Aakre. *Archaeological Fieldwork in Estonia*, 27–46.
- S122. Mäesalu, A. (1990). Sechs Holzkonstruktionen in Tartu (Lossi-Strasse). *Eesti Teaduste Akadeemia Toimetised. Ühiskonnateadused* 39(4), 446–452.
- S123. Mäesalu, A. (2001). Die Haustypen in hansezeitlichen Tartu (Dorpat). In: Lübecker Kolloquium zur Stadtarchäologie im Hanseraum III: der Hausbau/Bereich Archäologie der Hansestadt Lübeck; herausgegeben von Manfred Gläser. (Lübeck: Schmidt-Römhild), pp581–594.
- S124. Valk, H. (1999b). Aruanne arheoloogilistest kaevamistest Tartus Kuradisilla juures 1999. a. mais. (Excavation report, manuscript in the archaeology archive of the University of Tartu).
- S125. Tvauri, A. (2011b). Archaeological investigations at the courtyard of Jakobi Street 2/Lossi Street 3, Tartu. *Archaeological Fieldwork in Estonia 2010*, 179–186.
- S126. Malve, M., Roog, R., and Tvauri, A. (2012). Preliminary results of the rescue excavation in St Mary's churchyard and its surroundings in Tartu 2010–2011. *Archaeological Fieldwork in Estonia 2011*, 137–150.
- S127. Peets, J., Allmäe, R., and Maldre, L. (2011). Archaeological investigations of Pre-Viking Age burial boat in Salme village at Saaremaa. *Archeological Fieldwork in Estonia 2010*, 29–48.
- S128. Kustin, A. (1968). Aruanne Paatsa linnuse kaevamisest 1963. a. (Excavation report, manuscript in the archaeology archive of the University of Tartu).

- S129. Püüa, G., Toos, G., Alttoa, K., and Kadakas, V. (2013). Additions to the early construction history of the Kuressaare bishop's castle. *Archeological Fieldwork in Estonia* 2012, 171–184.
- S130. Lõugas, V. and Mägi-Lõugas, M. (1994). Excavations at Pöide stronghold have been finished. *Eesti NSV Teaduste Akadeemia Toimetised. Ühiskonnateadused* 43(4), 390–392.
- S131. Mägi, M. (2005). Viking Age harbour site at Tornimäe, eastern Saaremaa. *Archaeological Fieldwork in Estonia* 2004, 65–75.
- S132. Peets, J., Saage, R., and Maldre, L. (2012). The medieval and early modern smithy site of Käku. *Archaeological Fieldwork in Estonia*, 93–108.
- S133. Tvauri A. (2002). Aruanne arheoloogilistest kaevamistest Haapsalus Jaani 4 krundil 2002. aastal. (Excavation report, manuscript in the archaeology archive of the University of Tartu).
- S134. Lang V. (1988). Aruanne arheoloogilistest kaevamistest Iru linnuse keskvalil 1985. (ja 1986.a. Excavation report, manuscript in the archaeology archive of the University of Tartu).
- S135. Oras, E., Lang, V., Rannamäe, E., Varul, L., Konsa, M., Limbo-Simovart J., Vedru, G., Laneman, M., Malve, M., and Price, D. (2016). Tracing prehistoric migration: isotope analysis of Bronze and Pre-Roman Iron Age coastal burials in Estonia. *Estonian J. Archaeol.* 20(1), 3.
- S136. Lõugas, V. (1978). Archaeological research at Kaali meteorite crater. *Eesti NSV Teaduste Akadeemia Toimetised. Ühiskonnateadused* 27(1), 64–66.
- S137. Lõugas V. (1980). Arcahaeological excavations in the Kaali crater area. *Eesti NSV Teaduste Akadeemia Toimetised. Ühiskonnateadused* 29(4), 357–360.
- S138. Jonuks, T. (2012). Kiltsi mõis-keskaegsest kivimajast esindusliku härrastehooneni. *SA Virumaa Muuseumid toimetised*, 32–47.
- S139. Roog, R. and Malve, M. (2013). Rescue excavations on the settlement site and rural cemetery of Lohkva, Tartumaa. *Archaeological Fieldwork in Estonia* 2012, 241–250.
- S140. Tvauri, A. (2012). The Migration Period, Pre-Viking Age, and Viking Age in Estonia. *Estonian Archaeology* 4. (Tartu: Tartu University Press).
- S141. Lillak, A., and Valk, H. (2009b). Rescue Excavations on Alt-Laari Settlement Site, Tartumaa. *Archaeological Fieldwork in Estonia*, 65–71.
- S142. Valk, H. (1989b). Aruanne arheoloogilistest kaevamistest Sammaste kivikalmel 26.06.–31.07.1989. a. (Excavation report, manuscript in the archaeology archive of the University of Tartu).
- S143. Lehtosalo-Hilander, P.L. (1982). Luistari/I. The Graves. (Helsinki: Suomen muinaismuistoyhdistyksen aikakauskirja).
- S144. Tupala, U. (1999). Eläinuhreja vai teurasjätettä - Euran Luistarin rautakautisen ruumiskalmiston eläinluumateriaalin lähdekriittistä tarkastelua: University of Turku. (Department of Archaeology; MA thesis).
- S145. Naumann, E., Krzewinska, M., Götherström, A., and Gunilla, E. (2014). Slaves as burial gifts in Viking Age Norway? Evidence from stable isotope and ancient DNA analyses. *J. Archaeol. Sci.* 41, 533–540.

- S146. von Holstein, I.C., Walton Rogers, P., Craig, O.E., Penkman, K.E., Newton, J., and Collins, M.J. (2016). Provenancing Archaeological Wool Textiles from Medieval Northern Europe by Light Stable Isotope Analysis ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^2\text{H}$). *PloS one* 11, e0162330.
- S147. Goslar, T., Kalicki, T., Kaczanowska, M., Kozłowski, J.K. (2016). Stratigraphic sequence in trench A: complex II, layers 2–12 from the Early Neolithic to the Palaeolithic. In: Kaczanowska, M., Kozłowski, J.K., Sampson, A., eds. *The Sarakenos Cave at Akraephnion, Boeotia, Greece. Vol. II. The Early Neolithic, the Mesolithic and the Final Palaeolithic (Excavations in Trench A)*. (Kraków: The Polish Academy of Arts and Sciences), pp.18–33.
- S148. Niukkanen, M. (2002). *Sirpaleita Suurvalta-ajan Helsingistä*. (Hämeenlinna: Karisto Oy).
- S149. O'Shea, J.M. (1996). *The cemeteries of the Maros Group. Villagers of the Maros* (Springer), pp.55–80.
- S150. Sprincz, E., Beck, C.W. (1981). Classification of the amber beads of the Hungarian Bronze Age. *J. Field Archaeol.* 8, 469–485.
- S151. Zilhao, J. (2001). Radiocarbon evidence for maritime pioneer colonization at the origins of farming in west Mediterranean Europe. *P. Natl. Acad. Sci. U.S.A.* 98, 14180–14185.
- S152. Gabbianelli, F., Gargani, M., Pariset, L., Mariotti, M., Alhaique, F., De Minicis, E., Barelli, L., Ciammetti, E., Redi, F., and Valentini, A. (2015). Mitochondrial DNA analysis of medieval sheep (*Ovis aries*) in central Italy reveals the predominance of haplogroup B already in the Middle Ages. *Anim. Genet.* 46, 329–332.
- S153. Valenzuela-Lamas, S., Orengo, H.A., Bosch, D., Pellegrini, M., Halstead, P., Nieto-Espinet, A., Trentacoste, A., Jimenez-Manchon, S., Lopez-Reyes, D., and Jornet-Niella, R. (2018). Shipping amphorae and shipping sheep? Livestock mobility in the north-east Iberian peninsula during the Iron Age based on strontium isotopic analyses of sheep and goat tooth enamel. *PloS one* 13, e0205283.
- S154. Salzani, L. (2005). *La necropoli dell'età del bronzo all'Olmo di Nogara*. (Museo Civico di Storia Naturale, Milano).
- S155. Tafuri, M.A., Craig, O.E., Canci, A. (2009). Stable Isotope Evidence for the Consumption of Millet and Other Plants in Bronze Age Italy. *Amer. J. Phys. Anthropol.* 139, 146–153.
- S156. Baldoni, M., Scorrano, G., Gismondi, A., D'Agostino, A., Alexander, M., Gaspari, L., Vallelonga, F., Canini, A., Rickards, O., and Martinez-Labarga, C. (2018). Who were the miners of Allumiere? A multidisciplinary approach to reconstruct the osteobiography of an Italian worker community. *PloS one* 13, e0205362.
- S157. Gismondi, A., Baldoni, M., Gnes, M., Scorrano, G., D'Agostino, A., Di Marco, G., Calabria, G., Petrucci, M., Muldner, G., Von Tersch, M., et al. (2020). A multidisciplinary approach for investigating dietary and medicinal habits of the Medieval population of Santa Severa (7th–15th centuries, Rome, Italy). *PloS one* 15, e0227433.
- S158. Mirabaud, S., Rolando, C., and Regert, M. (2007). Molecular criteria for discriminating adipose fat and milk from different species by NanoESI MS and MS/MS of their triacylglycerols: application to archaeological remains. *Anal. Chem.* 79, 6182–6192.

- S159. Gustavsson, K. (1994). Franciskanerkloster på Kökar. Nytt ljus över medeltiden i Skärgårdshavet. *Historisk Tidskrift för Finland* 79, 494–518.
- S160. Caune, A. and Ose, I. (2004). Latvijas viduslaiku pilis IV. Latvijas 12. gadsimta beigās-17. gadsimta vācu piļu leksikons. (Rīga: Latvijas vēstures institūta apgāds).
- S161. Šnore E. (1978). Celtniecības liecības Kivtu apmetnē. In: *Arheoloģija un etnogrāfija XII. Apcerējumi par Latvijas teritorijas apmetnēm, celtnēm un tajās atrasto numismātisko materiālu pirmatnējās kopienas un feodālisma periodā.* (Rīga: Zinātne).
- S162. Atgāzis, M. (2006). Spietiņu uzkalniņkapi un to vieta Spietiņu un Plāteru senvietu kopā. In: Caune A, Mugarēvičs Ē, Ose I, Vasks A, editors. *Arheoloģija un etnogrāfija XXIII. Veltījums seno apgārbu pētniecei arheoloģei Dr. habil. hist. Annai Zariņai 85 gadu jubilejā.* (Rīga: Latvijas Vēstures Institūta Apgāds), pp.16–40.
- S163. Mugarēvičs, Ē. (1977). Oliņkalna un Lokstenes pilsnovadi. (Rīga: Zinātne).
- S164. Schultz, E.L. (1993). Mikkeli Moisio Latokallio/Moisioinpelto ja Mikkelin Mlk Kyyhkylä/Porrassalmenpelto: Finnish National Board of Antiquities. (Department of Archaeology).
- S165. Uotila K. (2011). Hautausmaan kaivaukset vuosina 2005–2007. In Naantalin luostarin rannassa. (Stranden vid Nådendals kloster. Edited by Uotila K. Eura: Kåkenhus-kirjat nro 3), pp.183–200.
- S166. Kallio, T. (2005). Oulu, Oulun kadut: Oikokatu, Kajaaninkatu, Torikatu, Ojakatu, Saaristonkatu, Franzenin puisto. Katutöiden arkeologinen valvonta: Finnish National Board of Antiquities. (Department of Monuments and Sites).
- S167. Hedman, S.D. and Olsen, B. (2009). Transition and order. A study of Sámi rectangular hearths in Pasvik, Arctic Norway. *Fennoscandia Archaeol.* 26, 3–23.
- S168. Oikarinen, T. (2009). Pietarsaari, 2/2/7-10, Lassfolkin kortteli (PSL-08). Kaupunkiarkeologinen pelastuskaivaus 19.5.-8.8.2008: Finnish National Board of Antiquities. (Department of Monuments and Sites).
- S169. Miettinen, M. (1993). Pihtiputaan Hämeensaari. Uutta tietoa Keski-Suomen historiasta. In Lapinraunioita ja Hiidenkiukaita. Volume 3. Edited by Purhonen P. Museovirasto: Arkeologian osasto, 52–64.
- S170. Wojtal, P., Makowiecki, D., Wertz, K., Wilczyński, J., Miękina, B., and Zabilska, M. (2010). Wstępne wyniki badań zooarcheologicznych szczątków kostnych z wykopalisk prowadzonych na Rynku Krakowskim w latach 2005–2007. In: Firlet, E., ed. (Krzysztofory. Zeszyty Naukowe Muzeum Historycznego Miasta Krakowa 28, część 2. Kraków: Muzeum Historyczne Miasta Krakowa), pp.137–150.
- S171. Vuorinen, J. (2009). Rakennukset ja rakentajat Raison lhalassa rautakauden lopulla ja varhaisella keskiajalla, Volume 281: University of Turku. (Annales Universitatis Turkuensis C).
- S172. Harlashov, B.N. (2014). = Харлашов БН. Отчет об археологических раскопках на территории объекта культурного наследия регионального значения "Селище 2-й пол. I тыс.н.э. в Выбутах" Псковского района Псковской области в 2013 г. Псков. (Excavation report, manuscript in the research archive of the Institute of Archaeology, Russian Academy of Science).

- S173. Ershova, E. (2015). Отчет об археологических раскопках на улице К.Маркса в г.Пскове в 2013 г. Новоторговский –XI раскоп. (Excavation report, manuscript in the research archive of the Institute of Archaeology, Russian Academy of Science).
- S174. Sedov. (1985). Средневековая Ладога. Новые археологические открытия и исследования. (Ленинград: Издательство “Наука” Ленинградское Отделение).
- S175. Poutiainen, H. (2000). Sysmä Ihananiemi. (Arkeologinen koekaivaus rautakautisella muinaisjäännösalueella: Lahti City Museum and Päijät-Häme Provincial Museum).
- S176. Achères ‘La Petite Ferme’. (2004). INRAP excavation. (unpublished).
- S177. Binois, A. (2015). Etude archéozoologique et paléopathologique des inhumations animales. In TeteghemCarlines 3. Rapport final d’opération. Ed M. Lançon. (INRAP Nord-Picardie).
- S178. Bazzanella, M. (2012). Italy: Neolithic and Bronze Age. In Gleba & Mannering, 203–214.
- S179. Bazzanella, M. and Mayr, A. (2009). I reperti tessili, le fusaiole e i pesi da telaio dalla palafitta di Molina di Ledro. (Trento: Provincia Autonoma di Trento).
- S180. Sabatini, Serena, Earle, T., and Cardarelli, A. (2018). Bronze Age Textile & Wool Economy: The Case of the Terramare Site of Montale, Italy. *Proc. Prehistoric Society* 84, 359–385.
- S181. Lopez-Costas, O. and Muldner, G. (2016). Fringes of the empire: Diet and cultural change at the Roman to post-Roman transition in NW Iberia. *Am. J. Phys. Anthropol.* 161(1), 141–154.
- S182. Gregoricka, L.A. (2013). Geographic origins and dietary transitions during the Bronze Age in the Oman Peninsula. *Am. J. Phys. Anthropol.* 152, 353–369.
- S183. Herva, V.P. (2002). Kaupunkiarkeologinen pelastuskaivaus, Tornion Keskikatu 29–35: University of Oulu. (Archaeological laboratory).
- S184. Riikonen, J. (2003). Gravundersökningar. Koroinen eläväksi-Korois till liv. Proceedings of the Korois-seminarium: 7–8 April 2001. Åbo landskapsmuseum rapporter 19, 21–24.
- S185. Alenius, T. and Laakso, V. (2006). Palaeoecology and archaeology of the village of Uukuniemi, Eastern Finland. *Acta. Borealia.* 23, 145–165.
- S186. Caffell, A. and Holst, M. (2011). Where Sky and Yorkshire and Water Meet: The Story of the Melton Landscape from Prehistory to the Present. Archaeological Excavations on the A63 Grade Separated Junction at Melton, East Yorkshire, Vol. Monograph 2 (ed. Chris Fenton–Thomas). (On–Site Archaeology, York).
- S187. Rusak, D.A., Marsico, R.M., and Taroli, B.L. (2011). Using Laser-Induced Breakdown Spectroscopy to Assess Preservation Quality of Archaeological Bones by Measurement of Calcium-to-Fluorine Ratios. *Appl. Spectrosc.* 65, 1193–1196.
- S188. Jones, J.R., Mulville, J.A., McGill, R.A., and Evershed, R.P. (2012). Palaeoenvironmental modelling of delta(13) C and delta(15) N values in the North Atlantic Islands: understanding past marine resource use. *Rapid Commun. Mass Sp.* 26, 2399–2406.
- S189. Salvagno, L., and Albarella, U. (2017). A morphometric system to distinguish sheep and goat postcranial

bones. PloS one 12, e0178543.

- S190. Ensminger, M.E. and Parker, R.O. (1986). *Sheep and Goat Science*, Fifth Edition. (Danville, Illinois: The Interstate Printers and Publishers Inc.).
- S191. Weaver, S. (2005). *Sheep: small-scale sheep keeping for pleasure and profit*. (Irvine, CA: Hobby Farm Press, an imprint of BowTie Press, a division of BowTie Inc.).
- S192. Ensminger, M.E. (1964). *Sheep and wool science*. (Interstate Printers & Publishers), p705.
- S193. Smith, B., Kennedy, G., Aseltine, M. (1997). *Beginning shepherd's manual*, Second Edition. (Ames, IA: Iowa State University Press).
- S194. Mair, V.H. (1998). *The Bronze Age and early Iron Age peoples of Eastern Central Asia*. (Institute for the Study of Man).
- S195. Chessa, B., Pereira, F., Arnaud, F., Amorim, A., Goyache, F., Mainland, I., Kao, R.R., Pemberton, J.M., Beraldi, D., Stear, M.J., et al. (2009). Revealing the history of sheep domestication using retrovirus integrations. *Science* 324, 532–536.
- S196. Green, M. (1980). Animal husbandry at Uruk in the Archaic period. *J. Near Eastern Studies* 39, 1–35.
- S197. Sherratt, A.G. (1981). Plough and pastoralism: aspects of the Secondary Products Revolution. In *Pattern of the Past* (eds I. Hodder, G. Isaac and N. Hammond). (Cambridge: Cambridge University Press), pp.261–306.
- S198. Good, I.L. (2001). Archaeological textiles: a review of current research. *Annu. Rev. Anthropol.* 30, 209–226.
- S199. Guo, W. (2012). *Archaeological study of late prehistoric society in xinjiang* (in Chinese). (Shanghai Classics Publishing House), p85.
- S200. Cottle David. (2010). *World Sheep and Wool Production*. (Nottingham University Press. Nottingham, United Kingdom).
- S201. Frei, K.M., Mannering, U., Kristiansen, K., Allentoft, M.E., Wilson, A.S., Skals, I., Tridico, S., Nosch, M.L., Willerslev, E., Clarke, L., et al. (2015). Tracing the dynamic life story of a Bronze Age Female. *Sci. Rep.* 5, 10431–10438.
- S202. Ryder, M.L. (1984). *Sheep*. In: Mason SL (ed) *Evolution of domesticated animals*. (London and New York: Longman).
- S203. Claude, A. (1994). The Archaeology of Africa. Food, Metals and Towns. (In: *Journal des africanistes*, tome 64, fascicule 2), pp.153–155.
- S204. Piao, W.Y. (2008). The origin and spread of silk tapestry (in Chinese). *Cultural relics appreciation and research*, 474–489.
- S205. Fernand, B. (1982). *The Wheels of Commerce*, vol 2 of *Civilization and Capitalism*. (New York: Harper & Row), pp.312–317
- S206. Skals, I. (2018). Wool textiles and archaeometry: testing reliability of archaeological wool fibre diameter measurements. *Danish J. Archaeol.* 7(2), 161–179.

- S207. Ryder, M.L. (1983). *Sheep and Man*. (London: Duckworth Press).
- S208. Davidson, A. (2006). *The Oxford Companion to Food*, T. Jaine, Ed. (Oxford University Press).
- S209. Feng, W.Q. (1991). A preliminary study on the formation of sheep breeds in ancient China (in Chinese). *Agr. Archaeol.* (3), 346–353.
- S210. Cixian cultural center. (1977). Dongwei tomb, dongchen village, cixian county, hebei (in Chinese). *Chinese Archaeology* (6), 33–42, 70, and 82–84.
- S211. Zhao, Y.X., Yang, J., Lv, F.H., Hu, X.J., Xie, X.L., Zhang, M., Li, W.R., Liu, M.J., Wang, Y.T., Li, J.Q., et al. (2017). Genomic Reconstruction of the History of Native Sheep Reveals the Peopling Patterns of Nomads and the Expansion of Early Pastoralism in East Asia. *Mol. Biol. Evol.* **34**, 2380–2395.
- S212. Kijas, J.W., Lenstra, J.A., Hayes, B., Boitard, S., Neto, L.R.P., San Cristobal, M., Servin, B., McCulloch, R., Whan, V., Gietzen, K., et al. (2012). Genome-wide analysis of the world's sheep breeds reveals high levels of historic mixture and strong recent selection. *Plos. Biol.* **10**, e1001258.
- S213. Meadows, J.R., Hanotte, O., Drogemuller, C., Calvo, J., Godfrey, R., Coltman, D., Maddox, J.F., Marzanov, N., Kantanen, J., and Kijas, J.W. (2006). Globally dispersed Y chromosomal haplotypes in wild and domestic sheep. *Anim. Genet.* **37**, 444–453.
- S214. Meadows, J.R., and Kijas, J.W. (2009). Re-sequencing regions of the ovine Y chromosome in domestic and wild sheep reveals novel paternal haplotypes. *Anim. Genet.* **40**, 119–123.
- S215. Ferencakovic, M., Curik, I., Perez-Pardal, L., Royo, L.J., Cubric-Curik, V., Fernandez, I., Alvarez, I., Kostelic, A., Sprem, N., Krapinec, K., et al. (2013). Mitochondrial DNA and Y-chromosome diversity in East Adriatic sheep. *Anim. Genet.* **44**, 184–192.
- S216. Zhang, M., Peng, W.F., Yang, G.L., Lv, F.H., Liu, M.J., Li, W.R., Liu, Y.G., Li, J.Q., Wang, F., Shen, Z.Q., et al. (2014). Y chromosome haplotype diversity of domestic sheep (*Ovis aries*) in northern Eurasia. *Anim. Genet.* **45**, 903–907.
- S217. Hu, X.J., Yang, J., Xie, X.L., Lv, F.H., Cao, Y.H., Li, W.R., Liu, M.J., Wang, Y.T., Li, J.Q., Liu, Y.G., et al. (2019). The Genome Landscape of Tibetan Sheep Reveals Adaptive Introgression from Argali and the History of Early Human Settlements on the Qinghai-Tibetan Plateau. *Mol. Biol. Evol.* **36**, 283–303.
- S218. Niemi, M., Blauer, A., Iso-Touru, T., Nystrom, V., Harjula, J., Taavitsainen, J.P., Stora, J., Liden, K., and Kantanen, J. (2013). Mitochondrial DNA and Y-chromosomal diversity in ancient populations of domestic sheep (*Ovis aries*) in Finland: comparison with contemporary sheep breeds. *Genet. Sel. Evol.* **45**, 2–15.
- S219. Li, X., Yang, J., Shen, M., Xie, X.L., Liu, G.J., Xu, Y.X., Lv, F.H., Yang, H., Yang, Y.L., Liu, C.B., et al. (2020). Whole-genome resequencing of wild and domestic sheep identifies genes associated with morphological and agronomic traits. *Nat. Commun.* **11**, 2815–2830.